Index of Authors

ABRAHAM, K. A. & EIKHOM, T. S. Determination of polyadenylate-rich ribonucleic acid in the nucleus and in the cytoplasm of plasmacytoma cells 669–674

AHLERS, J. The mechanism of hydrolysis of β-glycerophosphate by kidney alkaline phosphatase 535–546

AIROLDI, L. P. DA S. *see* DOONAN, S. 497–506

AITKEN, A. Prokaryote–eukaryote relationships and the amino acid sequence of plastocyanin from *Anabaena variabilis* 675–683

AKHTAR, M. *see* WILTON, D. C. 233–235

ANDANI, Z. *see* BROWN, C. R. 133–146, 147–154

ANDERSON, J. C. *see* LEE-OWN, V. 57–63

ANDREWS, P. T., JOHNSON, C. E., WALLBANK, B., CAMMACK, R., HALL, D. O. & RAO, K. K. X-ray photoelectron spectra of iron–sulphur proteins 471–474

ANOSIKE, E. O. & WATTS, D. C. Effects of anions on a monomeric and a dimeric arginine kinase 387–395

ANWAR, R. A. *see* GERBER, G. E. 685–695

ARCE, C. A. *see* RODRÍGUEZ, J. A. 115–121

ARCHIBALD, A. R. *see* HECKELS, J. E. 637–647

ASHTON, A. R. & POLYA, G. M. Higher-plant cyclic nucleotide phosphodiesterases. Resolution, partial purification and properties of three phosphodiesterases from potato tuber 329–339

ATKINS, G. L. & NIMMO, I. A. A comparison of seven methods for fitting the Michaelis–Menten equation 775–777

AVIRAM, I. *see* PETTIGREW, G. W. 155–167

AXELSSON, K. *see* MANNERVIK, B. 785–788

BACON, J. S. D., GORDON, A. H., MORRIS, E. J. & FARMER, V. C. Acetyl groups in cell-wall preparations from higher plants 485–487

BADDILEY, J. *see* HECKELS, J. E. 637–647; LAMBERT, P. A. 519–524

BAILEY, A. J. *see* ROBINS, S. P. 381–385

BARDSLEY, W. G. & CHILDS, R. E. Sigmoid curves, non-linear double-reciprocal plots and allosterism 313–328

BARENHOLZ, Y. *see* QUINN, P. J. 199–208

BARON, A. J. *see* HOLDER, A. A. 757–773; WOOTTON, J. C. 749–754

BARRA, D. *see* DOONAN, S. 497–506

BARRA, H. S. *see* RODRÍGUEZ, J. A. 115–121

BARUA, N. N. *see* FITT, P. S. 719–724

BAXTER, E. & MUIR, H. The nature of the protein moieties of cartilage proteoglycans of pig and ox 657–668

BAYLEY, P. M. *see* THORNTON, J. M. 585–596

BLACKBURN, G. M. & TAUSSIG, P. E. The photocarcinogenicity of anthracene: photochemical binding to deoxyribonucleic acid in tissue culture 289–291

BOSSA, F. *see* DOONAN, S. 497–506

BOTTOMLEY, R. C. & TRAYER, I. P. Affinity chromatography of immobilized actin and myosin 365–379

BREMNER, I. & DAVIES, N. T. The induction of metallothionein in rat liver by zinc injection and restriction of food intake 733–738

BRISTOW, A. F. *see* VIRDEN, R. 397–401

BRITTAIN, T. & GREENWOOD, C. Kinetic studies on [*N*-formyltryptophyl]cytochrome *c* 713–717

BRITTAIN, T. & GREENWOOD, C. Kinetic studies on mammalian cytochrome *c* modified with 2-hydroxy-5-nitrobenzyl bromide 179–185

BROWN, C. R., ANDANI, Z. & HARTREE, E. F. Studies on ram acrosin. Fluorimetric titration of operational molarity with 4-methylumbelliferyl *p*-guanidinobenzoate 147–154

BROWN, C. R., ANDANI, Z. & HARTREE, E. F. Studies on ram acrosin. Isolation from spermatozoa, activation by cations and organic solvents, and influence of cations on its reaction with inhibitors 133–146

BURNS, G. R. J. & WYNN, C. H. Studies on the arylsulphatase and phenol sulphotransferase activities of *Aspergillus oryzae* 697–705

BUTTERWORTH, R. M., GIBSON, J. F. & WILLIAMS, J. Electron-paramagnetic-resonance spectroscopy of iron-binding fragments of hen ovotransferrins 559–563

CAMMACK, R. *see* ANDREWS, P. T. 471–474

CAPUTTO, R. *see* RODRÍGUEZ, J. A. 115–121

CARLONI, M. *see* DOONAN, S. 497–506

CARRELL, R. W. *see* CHUA, C. G. 259–269

CHAMBERS, G. K. *see* HOLDER, A. A. 757–773; WOOTTON, J. C. 739–748

CHAPMAN, M. J., MILLS, G. L. & LEDFORD, J. H. The distribution and partial characterization of the serum apolipoproteins in the guinea pig 423–436

CHEN, S.-S. & ENGEL, P. C. Horse liver alcohol dehydrogenase. A study of the essential lysine residue 627–635

CHEN, S.-S. & ENGEL, P. C. Modification of pig M_4 lactate dehydrogenase by pyridoxal 5′-phosphate Demonstration of an essential lysine residue 107–113

CHEN, S.-S. & ENGEL, P. C. Ox liver glutamate dehydrogenase. The role of lysine-126 reappraised in the light of studies of inhibition and inactivation by pyridoxal 5′-phosphate 619–626

CHENG, K.-W. Properties of follicle-stimulating-hormone receptor in cell membranes of bovine testis 123–132

CHILDS, R. E. *see* BARDSLEY, W. G. 313–238

CHIU, A. T., RYAN, J. W., RYAN, U. S. & DORER, F. E. A sensitive radiochemical assay for angiotensin-converting enzyme (kinase II) 297–300

CHUA, C. G., CARRELL, R. W. & HOWARD, B. H. The amino acid sequence of the α chain of the major haemoglobin of the rat (*Rattus norvegicus*) 259–269

COLE, H. A. & PERRY, S. V. The phosphorylation of troponin I from cardiac muscle 525–533

CORNISH-BOWDEN, A. The use of the direct linear plot for determining initial velocities 305–312

CORRAN, P. H. & WALEY, S. G. The reaction of penicillin with proteins 357–364

DAVIES, D. D. *see* DO NASCIMENTO, K. H. 349–355, 553–557

DAVIES, N. T. *see* BREMNER, I. 733–738

DAWSON, R. M. The effect of some amine oxides and disulphide compounds on the activity of acetylcholinesterase 293–295

DELL, A. *see* MORRIS, H. R. 754–755

DEUMLING, B. *see* SINCLAIR, J. 31–38

DEVONSHIRE, A. L. Studies of the acetylcholinesterase from houseflies (*Musca domestica* L.) resistant and susceptible to organophosphorus insecticides 463–469
DO NASCIMENTO, K. H. & DAVIES, D. D. The stereospecificity of sequential nicotinamide–adenine dinucleotide-dependent oxidoreductases in relation to the evolution of metabolic sequences 553–557
DO NASCIMENTO, K. H., DAVIES, D. D. & PATIL, K. D. Undirectional inhibition and activation of 'malic' enzyme of *Solanum tuberosum* by *meso*-tartrate 349–355
DONDI, P. G. Analysis, by two-dimensional gel electrophoresis, of ribosome crystal proteins 475–476
DOONAN, H. J. *see* DOONAN, S. 497–506
DOONAN, S., DOONAN, H. J., HANFORD, R., VERNON, C. A., WALKER, J. M., AIROLDI, L. P. DA S., BOSSA, F., BARRA, D., CARLONI, M., FASELLA, P. & RIVA, F. The primary structure of aspartate aminotransferase from pig heart muscle. Digestion with a proteinase having specificity for lysine residues 497–506
DORER, F. E. *see* CHIU, A. T. 297–300
DOWER, S. K., DWEK, R. A., MCLAUGHLIN, A. C., MOLE, L. E., PRESS, E. M. & SUNDERLAND, C. A. The binding of lanthanides to non-immune rabbit immunoglobulin G and its fragments 73–82
DREWRY, D. T., SYMES, K. C., GRAY, G. W. & WILKINSON, S. G. Studies of polysaccharide fractions from the lipopolysaccharide of *Pseudomonas aeruginosa* N.C.T.C. 1999 93–106
DUMONT, J. E. *see* SWILLENS, S. 779–782
DUNKERTON, J. & JAMES, S. P. Purification of 2-oxoaldehyde dehydrogenase and its dependence on unusual amines 609–617
DWEK, R. A. *see* DOWER, S. K. 73–82

EAGLES, P. A. M. *see* PAU, R. N. 707–712
EGGITT, M. J. & SCRAGG, A. H. Messenger activity of ribonucleic acid from yeast mitochondria 507–512
EHNHOLM, C., KINNUNEN, P. K. J., HUTTUNEN, J. K., NIKKILÄ, E. A. & OHTA, M. Purification and characterization of lipoprotein lipase from pig myocardium 649–655
EIKHOM, T. S. *see* ABRAHAM, K. A. 669–674
ENGEL, P. C. *see* CHEN, S.-S. 107–113, 619–626, 627–635

FARMER, V. C. *see* BACON, J. S. D. 485–487
FASELLA, P. *see* DOONAN, S. 497–506
FINCHAM, J. R. S. *see* HOLDER, A. A. 757–773; WOOTTON, J. C. 739–748, 749–754
FITT, P. S., PETERKIN, P. I. & BARUA, N. N. The relationship between the deoxyribonucleic acid-bound and low-molecular-weight soluble forms of *Halobacterium* cutirubrum deoxyribonucleic acid-dependent ribonucleic acid polymerase 719–724
FITZGERALD, J. W. Secondary alkylsulphatases in a strain of *Comamonas terrigena* 477–480

GARRICK, L. M., SHARMA, V. S., MCDONALD, M. J. & RANNEY, H. M. Rat haemoglobin heterogeneity. Two structurally distinct α chains and functional behaviour of selected components 245–258
GAYATHRI DEVI, B., SHAILA, M. S., RAMAKRISHNAN, T. & GOPINATHAN, K. P. The purification and properties of peroxidase in *Mycobacterium tuberculosis* H37Rv and its possible role in the mechanism of action of isonicotinic acid hydrazide 187–197
GERBER, G. E. & ANWAR, R. A. Comparative studies of the cross-linked regions of elastin from bovine ligamentum nuchae and bovine, porcine and human aorta 685–695
GIBSON, J. F. *see* BUTTERWORTH, R. M. 559–563
GOOCH, J. & HAWTREY, A. O. Synthesis of thiol-containing analogues of puromycin and a study of their interaction with *N*-acetylphenylalanyl-transfer ribonucleic acid on ribosomes to form thioesters 209–220
GOPINATHAN, K. P. *see* GAYATHRI DEVI, B. 187–197
GORDON, A. H. *see* BACON, J. S. D. 485–487
GRAND, R. J. A. *see* WILKINSON, J. M. 493–496
GRAY, G. W. *see* DREWRY, D. T. 93–106
GREENWOOD, C. *see* BRITTAIN, T. 179–185, 713–717; MOORE, T. A. 169–177
GREGORY, R. P. F. *see* SCOTT, B. 341–347
GRIENENBERGER, J. M. & SIMON, D. Structure and biosynthesis of the ribosomal ribonucleic acids from the oncogenic bacterium *Agrobacterium tumefaciens* 23–30

HALFORD, S. E. Stopped-flow fluorescence studies on saccharide binding to lysozyme 411–422
HALL, D. O. *see* ANDREWS, P. T. 471–474
HALLAK, M. E. *see* RODRÍGUEZ, J. A. 115–121
HANCOCK, I. C. *see* LAMBERT, P. A. 519–524
HANFORD, R. *see* DOONAN, S. 497–506
HARTREE, E. F. *see* BROWN, C. R. 133–146, 147–154
HAWTREY, A. O. *see* GOOCH, J. 209–220
HECKELS, J. E., ARCHIBALD, A. R. & BADDILEY, J. Studies on the linkage between teichoic acid and peptidoglycan in a bacteriophage-resistant mutant of *Staphylococcus aureus* H 637–647
HOLBROOK, I. B. & MOLAN, P. C. The identification of a peptide in human parotid saliva particularly active in enhancing the glycolytic activity of the salivary microorganisms 489–492
HOLDER, A. A., WOOTTON, J. C., BARON, A. J., CHAMBERS, G. K. & FINCHAM, J. R. S. The amino acid sequence of *Neurospora* NADP-specific glutamate dehydrogenase. Peptic and chymotryptic peptides and the complete sequence 757–773
HOLLAWAY, M. R. & WHITE, H. A. A double-beam rapid-scanning stopped-flow spectrophotometer 221–231
HÖÖK, M. *see* JANSSON, L. 49–55
HOWARD, B. H. *see* CHUA, C. G. 259–269
HUTTUNEN, J. K. *see* EHNHOLM, C. 649–655

INGLE, J. *see* SINCLAIR, J. 31–38
ISHIBASHI, S. *see* OUCHI, M. 481–483
IUPAC COMMISSION ON THE NOMENCLATURE OF ORGANIC CHEMISTRY & IUPAC–IUB COMMISSION ON BIOCHEMICAL NOMENCLATURE Nomenclature of α-amino acids (Recommendations, 1974) 1–16
IUPAC–IUB COMMISSION ON BIOCHEMICAL NOMENCLATURE *see* IUPAC COMMISSION ON THE NOMENCLATURE OF ORGANIC CHEMISTRY 1–16

JACKSON, A. A. *see* WOOTTON, J. C. 739–748
JAMES, S. P. *see* DUNKERTON, J. 609–617

JANSSON, L., HÖÖK, M., WASTESON, Å. & LINDAHL, U. Biosynthesis of heparin. Solubilization and partial characterization of *N*- and *O*-sulphotransferases 49–55
JOHNSON, C. E. *see* ANDREWS, P. T. 471–474

KALGHATGI, K. K. & SUBBA RAO, P. V. Microbial L-phenylalanine ammonia-lyase. Purification, subunit structure and kinetic properties of the enzyme from *Rhizoctonia solani* 65–72
KALYANARAMAN, V. S., KUMAR, S. A. & MAHADEVAN, S. Oxidase–peroxidase enzymes of *Datura innoxia*. Oxidation of reduced nicotinamide–adenine dinucleotide in the presence of formylphenylacetic acid ethyl ester 577–584
KALYANARAMAN, V. S., MAHADEVAN, S. & KUMAR, S. A. Oxidase–peroxidase enzymes of *Datura innoxia*. Oxidation of formylphenylacetic acid ethyl ester 565–576
KINNUNEN, P. K. J. *see* EHNHOLM, C. 649–655
KUMAR, S. A. *see* KALYANARAMAN, V. S. 565–576, 577–584
KVAMME, E. & TORGNER, I. AA. Regulatory effects of fatty acyl-coenzyme A derivatives on phosphate-activated pig brain and kidney glutaminase *in vitro* 83–91

LAMBERT, P. A., HANCOCK, I. C. & BADDILEY, J. The interaction of magnesium ions with teichoic acid 519–524
LAYCOCK, M. V. The amino acid sequence of cytochrome *f* from the brown alga *Alaria esculenta* (L.) Grev. 271–279
LEA, P. J. *see* MIFLIN, B. J. 403–409
LEDFORD, J. H. *see* CHAPMAN, M. J. 423–436
LEE-OWN, V. & ANDERSON, J. C. The isolation of collagen-associated proteoglycan from bovine nasal cartilage and its preferential interaction with α_2 chains of type I collagen 57–63
LINDAHL, U. *see* JANSSON, L. 49–55
LOENING, U. E. *see* SCHUCH, W. 17–22
LUCY, J. A. *see* MAGGIO, B. 597–608

MAGGIO, B. & LUCY, J. A. Studies on mixed monolayers of phospholipids and fusogenic lipids 597–608
MAHADEVAN, S. *see* KALYANARAMAN, V. S. 565–576, 577–584
MANNERVIK, B. & AXELSSON, K. Reduction of disulphide bonds in proteins and protein mixed disulphides by a thioltransferase in rat liver cytosol 785–788
MCDONALD, M. J. *see* GARRICK, L. M. 245–258
MCINTYRE, R. J. & VAUGHAN, P. F. T. Kinetic studies on the hydroxylation of *p*-coumaric acid to caffeic acid by spinach-beet phenolase 447–461
MCLAUGHLIN, A. C. *see* DOWER, S. K. 73–82
MIFLIN, B. J. & LEA, P. J. Glutamine and asparagine as nitrogen donors for reductant-dependent glutamate synthesis in pea roots 403–409
MILLS, G. L. *see* CHAPMAN, M. J. 423–436
MOLAN, P. C. *see* HOLBROOK, I. B. 489–492
MOLE, L. E. *see* DOWER, S. K. 73–82
MOORE, T. A. & GREENWOOD, C. A method for investigating the effect of temperature on the 695 nm band of insoluble cytochrome *c* 169–177
MORRIS, E. J. *see* BACON, J. S. D. 485–487
MORRIS, H. R. & DELL, A. The sequence of the blocked *N*-terminal peptide from *Neurospora* glutamate dehydrogenase 754–755
MOSBY, L. J. *see* WOOD, E. J. 437–445
MUIR, H. *see* BAXTER, E. 657–668

NIKKILÄ, E. A. *see* EHNHOLM, C. 649–655
NIMMO, I. A. *see* ATKINS, G. L. 755–777

OHTA, M. *see* EHNHOLM, C. 649–655
ONAJOBI, F. D. Effects of adenine nucleotides on rice-root adenosine triphosphate sulphurylase activity *in vitro* 301–304
OUCHI, M. & ISHIBASHI, S. Discrimination of antigenic site and thiol-inhibitor-sensitive site of hexokinase isoenzymes 481–483

PAIN, R. H. *see* VIRDEN, R. 397–401
PATIL, K. D. *see* DO NASCIMENTO, K. H. 349–345
PAU, R. N. & EAGLES, P. A. M. The isolation of an *o*-diphenol oxidase from third-instar larvae of the blowfly *Calliphora erythrocephala* 707–712
PERRY, S. V. *see* COLE, H. A. 525–533
PETERKIN, P. I. *see* FITT, P. S. 719–724
PETTIGREW, G. W., AVIRAM, I. & SCHEJTER, A. Physiochemical properties of two atypical cytochromes *c*, *Crithidia* cytochrome *c*-557 and *Euglena* cytochrome *c*-558 155–167
POLYA, G. M. *see* ASHTON, A. R. 329–339
PRESS, E. M. Fixation of the first component of complement by immune complexes: effect of reduction and fragmentation of antibody 285–288
PRESS, E. M. *see also* DOWER, S. K. 73–82

QUINN, P. J. & BARENHOLZ, Y. A comparison of the activity of phosphatidylinositol phosphodiesterase against substrate in dispersions and as monolayers at the air–water interface 199–208

RAMAKRISHNAN, T. *see* GAYATHRI DEVI, B. 187–197
RANNEY, H. M. *see* GARRICK, L. M. 245–258
RAO, K. K. *see* ANDREWS, P. T. 471–474
REDMAN, D. G. Structural studies on wheat (*Triticum aestivum*) proteins lacking phenylalanine and histidine residues 725–732
RIVA, F. *see* DOONAN, S. 497–506
ROBINS, S. P. & BAILEY, A. J. The chemistry of the collagen cross-links. The mechanism of stabilization of the reducible intermediate cross-links 381–385
RODRÍGUEZ, J. A., BARRA, H. S., ARCE, C. A., HALLAK, M. E. & CAPUTTO, R. The reciprocal exclusion by L-dopa (L-3,4-dihydroxyphenylalanine) and L-tyrosine of their incorporation as single units into a soluble rat brain protein 115–121
RYAN, J. W. *see* CHIU, A. T. 297–300
RYAN, U. S. *see* CHIU, A. T. 297–300

SCHEJTER, A. *see* PETTIGREW, G. W. 155–167
SCHUCH, W. & LOENING, U. E. The ribosomal ribonucleic acid of *Agrobacterium tumefaciens* 17–22
SCOTT, B. & GREGORY, R. P. F. Properties of protein–chlorophyll complexes from pea (*Pisum sativum L.*) leaves. The organization of chlorophyll 341–347
SCRAGG, A. H. *see* EGGITT, M. J. 507–512

SERAFINI-CESSI, F. Determination of 2-Acetamido-2-deoxy-D-galactose and mechanism of formation of chromogens 513–517
SHAILA, M. S. *see* GAYATHRI DEVI, B. 187–197
SHARMA, V. S. *see* GARRICK, L. M. 245–258
SIMON, D. *see* GRIENENBERGER, J. M. 23–30
SINCLAIR, J., WELLS, R., DEUMLING, B. & INGLE, J. The complexity of satellite deoxyribonucleic acid in a higher plant 31–38
STEINERT, P. M. The extraction and characterization of bovine epidermal α-keratin 39–48
SUBBA RAO, P. V. *see* KALGHATGI, K. K. 65–72
SUNDERLAND, C. A. *see* DOWER, S. K. 73–82
SWILLENS, S. & DUMONT, J. E. A pitfall in the interpretation of data on ligand–protein interaction 779–782
SYMES, K. C. *see* DREWRY, D. T. 93–106

TAUSSIG, P. E. *see* BLACKBURN, G. M. 289–291
TAYLOR, J. G. *see* WOOTTON, J. C. 739–748
THORNTON, J. M. & BAYLEY, P. M. Conformational energy calculations for dinucleotide molecules. A study of the component mononucleotides adenosine 5′-monophosphate, nicotinamide mononucleotide and adenosine 3′-monophosphate 585–596
TORGNER, I. AA. *see* KVAMME, E. 83–91
TRAYER, I. P. *see* BOTTOMLEY, R. C. 365–379

VAUGHAN, P. F. T. *see* MCINTYRE, R. J. 447–461
VERNON, C. A. *see* DOONAN, S. 497–506
VIRDEN, R., BRISTOW, A. F. & PAIN, R. H. The active site of penicillinase from *Staphylococcus aureus* PC1. Isolation of a specific covalent complex with the substrate quinacillin 397–401

WALEY, S. G. The pH-dependence and group modification of β-lactamase I 547–551
WALEY, S. G. *see also* CORRAN, P. H. 357–364
WALKER, J. M. *see* DOONAN, S. 497–506
WALLBANK, B. *see* ANDREWS, P. T. 471–474
WASTESON, Å. *see* JANSSON, L. 49–55
WATTS, D. C. *see* ANOSIKE, E. O. 387–395
WEITZMAN, P. D. J. Evidence against the proposed interaction of thionitrobenzoate with protein disulphide bonds 281–283
WELBOURN, A. P. *see* WILKINSON, S. G. 783–784
WELLS, R. *see* SINCLAIR, J. 31–38
WHITE, H. A. *see* HOLLOWAY, M. R. 221–231
WILKINSON, J. M. & GRAND, R. J. A. The amino acid sequence of troponin I from rabbit skeletal muscle 493–496
WILKINSON, S. G. & WELBOURN, A. P. 2-Amino-2-deoxygalacturonic acid in lipopolysaccharides from *Pseudomonas aeruginosa* 783–784
WILKINSON, S. G. *see also* DREWRY, D. T. 93–106
WILLIAMS, J. Iron-binding fragments from the carboxyl-terminal region of hen ovotransferrin 237–244
WILLIAMS, J. *see also* BUTTERWORTH, R. M. 559–563,
WILTON, D. C. & AKHTAR, M. The mechanism of C-4 demethylation during cholesterol biosynthesis. Evidence for a decarboxylation mechanism not involving a Schiff-base intermediate 233–235
WOOD, E. J. & MOSBY, L. J. Physicochemical properties of *Planorbis corneus* erythrocruorin 437–445
WOOTTON, J. C., BARON, A. J. & FINCHAM, J. R. S. The amino acid sequence of *Neurospora* NADP-specific glutamate dehydrogenase. Peptides from digestion with a staphylococcal proteinase 749–754
WOOTTON, J. C., TAYLOR, J. G., JACKSON, A. A., CHAMBERS, G. K. & FINCHAM, J. R. S. The amino acid sequence of *Neurospora* NADP-specific glutamate dehydrogenase. The tryptic peptides 739–748
WOOTTON, J. C. *see also* HOLDER, A. A. 757–773
WYNN, C. H. *see* BURNS, G. R. J. 697–705

Index of Subjects

2-Acetamido-2-deoxy-D-galactose, *see* *N*-Acetyl-D-galactosamine

Acetate ions, effects of, and other anions on the activity of monomeric (lobster) and dimeric (sea-cucumber) arginine kinases (Anosike, E. O. & Watts, D. C.) 387–395

Acetylcholinesterase, erythrocyte, ox, effects of some amine oxides and disulphide compounds on the activity of (Dawson, R. M.) 293–295

Acetylcholinesterase, properties of, from houseflies resistant and susceptible to organophosphorus insecticides (Devonshire, A. L.) 463–469

N-Acetyl-D-galactosamine, determination of, and the mechanism of the formation of chromogens (Serafini-Cessi, F.) 513–517

N-Acetyl-D-glucosamine, stopped-flow fluorescence studies on the binding of the β-(1 → 4)-linked trimer of, to hen's-egg-white lysozyme (Halford, S. E.) 411–422

Acetyl groups, detection of, in preparations of cell walls from grasses and other higher plants (Bacon, J. S. D., Gordon, A. H., Morris, E. J. & Farmer, V. C.) 485–487

N-Acetylhexosamines, determination of, and the mechanism of the formation of chromogens (Serafini-Cessi, F) 513–517

N-Acetylphenylalanyl-transfer ribonucleic acid, interaction of thiol-containing analogues of puromycin with, on ribosomes from rat liver and *Escherichia coli* to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220

Acrosin, partial purification and properties of, from sheep spermatozoa (Brown, C. R., Andani, Z. & Hartree, E. F.) 133–146

Acrosin, spermatozoal, sheep, determination of the operational molarity of, by fluorimetric titration with 4-methylumbelliferyl *p*-guanidinobenzoate (Brown, C. R., Andani, Z. & Hartree, E. F.) 147–154

Actin, skeletal-muscle, rabbit, use of affinity chromatography for assessment of the interactions of, and myosin (Bottomley, R. C. & Trayer, I. P.) 365–379

Acyl-coenzyme A, regulatory effects of, on the activity of pig brain and kidney phosphate-activated glutaminase *in vitro* (Kvamme, E. & Torgner, I. Aa.) 83–91

Adenine nucleotides, effects of, on the activity of rice-root adenosine triphosphate sulphurylase *in vitro* (Onajobi, F. D.) 301–304

Adenosine 3′-monophosphate, calculations of the conformational energy of, and other mononucleotide moieties in dinucleotides (Thornton, J. M. & Bayley, P. M.) 585–596

Adenosine 5′-monophosphate, calculations of the conformational energy of, and other mononucleotide moieties in dinucleotides (Thornton, J. M. & Bayley, P. M.) 585–596

Adenosine triphosphate sulphurylase, rice-root, effects of adenine nucleotides on the activity of, *in vitro* (Onajobi, F. D.) 301–304

Aerobacter aerogenes N.C.I.B. 8017, stereospecificity of nicotinamide–adenine dinucleotide-dependent mannitol 1-phosphate dehydrogenase and sorbitol 6-phosphate dehydrogenase from (do Nascimento, K. H. & Davies, D. D.) 553–557

Agrobacterium tumefaciens, presence of nicks in the 23S ribosomal ribonucleic acid of (Schuch, W. & Loening, U. E.) 17–22

Agrobacterium tumefaciens, structure and biosynthesis of the ribosomal ribonucleic acids of (Grienenberger, J. M. & Simon, D.) 23–30

Alaria esculenta, amino acid sequence of cytochrome *f* from (Laycock, M. V.) 271–279

Alcohol dehydrogenase, liver, horse, inactivation of, by reaction of pyridoxal 5′-phosphate with the essential lysine residue (Chen, S.-S. & Engel, P. C.) 627–635)

Alcohol dehydrogenase, liver, horse, use of a double-beam rapid-scanning stopped-flow spectrophotometer to measure rapid changes in the spectrum of reduced nicotinamide–adenine dinucleotide during the reaction catalysed by (Hollaway, M. R. & White, H. A.) 221–231

Alkaline phosphatase, *see* Phosphatase, alkaline

Alkylsulphatases, secondary, occurrence of, in a strain of *Comamonas terrigena* (Fitzgerald, J. W.) 477–480

Amine oxides, effects of various disulphides and, on the activity of ox erythrocyte acetylcholinesterase (Dawson, R. M.) 293–295

Amines, dependence on, of the activity of sheep liver 2-oxo aldehyde dehydrogenase (Dunkerton, J. & James, S. P.) 609–617

Amino acid compositions of three wheat-flour proteins that lack phenylalanine and histidine residues (Redman, D. G.) 725–732

Amino acid sequence of *Alaria esculenta* cytochrome *f* (Laycock, M. V.) 271–279

Amino acid sequence of *Anabaena variabilis* plastocyanin and its comparison with those of plastocyanins from *Chlorella fusca* and higher plants (Aitken, A.) 675–683

Amino acid sequence of *Neurospora crassa* nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase (Holder, A. A., Wootton, J. C., Baron, A. J., Chambers, G. K., & Fincham, J. R. S.) 757–773

Amino acid sequence of rabbit skeletal-muscle troponin I (Wilkinson, J. M. & Grand, R. J. A.) 493–496

Amino acid sequence of the blocked *N*-terminal peptide of *Neurospora crassa* nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase (Morris H. R. & Dell, A.) 754–755

Amino acid sequence, partial, of the α chain of the major rat haemoglobin (Chua, C. G., Carrell, R. W. & Howard, B. H.) 259–269

Amino acid sequence, use of digestion with *Armillaria mellea* proteinase having specificity for lysine residues in the establishment of, of pig heart-muscle aspartate aminotransferase (Doonan, S., Doonan, H. J., Hanford, R., Vernon, C. A., Walker, J. M., Airoldi, L. P. da S., Bossa, F., Barra, D., Carloni, M., Fasella, P. & Riva, F.) 497–506

Amino acid sequences of peptides obtained by the digestion of *Neurospora crassa* nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase with a staphylococcal proteinase (Wootton, J. C., Baron, A. J. & Fincham, J. R. S.) 749–754

Amino acid sequences of peptides obtained by the digestion of *Neurospora crassa* nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase with pepsin and chymotrypsin (Holder, A. A., Wootton, J. C., Baron, A. J., Chambers, G. K. & Fincham, J. R. S.) 757–773

Amino acid sequences of peptides obtained by the digestion of *Neurospora crassa* nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase with trypsin (Wootton, J. C., Taylor, J. G., Jackson, A. A., Chambers, G. K. & Fincham, J. R. S.) 739–748

Amino acid sequences, partial, of two structurally distinct α chains of rat haemoglobins (Garrick, L. M., Sharma, V. S., McDonald, M. J. & Ranney, H. M.) 245–258

α-Amino acids, nomenclature of (IUPAC Commission on the Nomenclature of Organic Chemistry & IUPAC–IUB Commission on Biochemical Nomenclature) 1–16

2-Amino-2-deoxygalacturonic acid, identification of, as a component of *Pseudomonas aeruginosa* N.C.T.C. 8505 lipopolysaccharide (Wilkinson, S. G. & Welbourn, A. P.) 783–784

2-Aminopropan-1-ol, dependence of the activity of sheep liver 2-oxo aldehyde dehydrogenase on, and other amines (Dunkerton, J. & James, S. P.) 609–617

Ampicillin, dependence on pH of the hydrolysis of benzylpenicillin and, by *Bacillus cereus* 569/H β-lactamase I (Waley, S. G.) 547–551

Anabaena variabilis, amino acid sequence of plastocyanin from, and its comparison with those of plastocyanins from *Chlorella fusca* and higher plants (Aitken, A.) 675–683

Angiotensin-converting enzyme (kinase II), lung, sensitive radiochemical method for assay of the activity of (Chiu, A. T., Ryan, J. W., Ryan, U. S. & Dorrer, F. E.) 297–300

Anions, effects of, on the activity of monomeric (lobster) and dimeric (sea-cucumber) arginine kinases (Anosike, E. O. & Watts, D. C.) 387–395

Anthracene, photochemical binding of, to deoxyribonucleic acid of monkey kidney epithelial cells and human skin epithelial cells in tissue culture (Blackburn, G. M. & Taussig, P. E.) 289–291

Antigen, O, characterization of polysaccharide fractions obtained by mild acid hydrolysis of *Pseudomonas aeruginosa* N.C.T.C. 1999 lipopolysaccharide active as (Drewry, D. T., Symes, K. C., Gray, G. W. & Wilkinson, S. G.) 93–106

Aorta, ox, pig and human, comparative studies of the cross-linked regions of elastins from, and ox ligamentum nuchae (Gerber, G. E. & Anwar, R. A.) 685–695

Apolipoproteins, serum, guinea-pig, distribution and partial characterization of (Chapman, M. J., Mills, G. L. & Ledford, J. H.) 423–436

Arginine kinases, monomeric (lobster) and dimeric (sea-cucumber), effects of anions on the activity of (Anosike, E. O. & Watts, D. C.) 387–395

Armillaria mellea, use of digestion with proteinase from, having specificity for lysine residues in the establishment of the amino acid sequence of pig heart-muscle aspartate aminotransferase (Doonan, S., Doonan, H. J., Hanford, R., Vernon, C. A., Walker, J. M., Airoldi, L. P. da S., Bossa, F., Barra, D., Carloni, M., Fasella, P. &. & Riva, F.) 497–506

Arylsulphatases, *Aspergillus oryzae*, properties of, one possessing phenol sulphotransferase activity (Burns, G. R. J. & Wynn, C. H.) 697–705

Asparagine, glutamine and, as nitrogen donors for the reductant-dependent biosynthesis of glutamate in pea roots (Miflin, B. J. & Lea, P. J.) 403–409

Aspartate aminotransferase, heart-muscle, pig, use of digestion with *Armillaria mellea* proteinase having specificity for lysine residues in the establishment of the amino acid sequence of (Doonan, S., Doonan, H. J., Hanford, R., Vernon, C. A., Walker, J. M., Airoldi, L. P. da S., Bossa, F., Barra, D., Carloni, M., Fasella, P. & Riva, F.) 497–506

Aspergillus oryzae, properties of arylsulphatases, one possessing phenol sulphotransferase activity, from (Burns, G. R. J. & Wynn, C. H.) 697–705

Bacillus cereus 569/H, dependence on pH of the hydrolysis of benzylpenicillin and ampicillin by β-lactamase I from (Waley, S. G.) 547–551

Beet (*Beta vulgaris*) leaves, kinetics of the hydroxylation of *p*-coumarate to caffeate by phenolase from (McIntrye, R. J. & Vaughan, P. F. T.) 447–461

Benzylpenicillin, dependence on pH of the hydrolysis of ampicillin and, by *Bacillus cereus* 569/H β-lactamase I (Waley, S. G.) 547–551

Benzylpenicillin, reaction of, with pig insulin and with hen's-egg-white lysozyme (Corran, P. H. & Waley, S. G.) 357–364

Beta vulgaris, *see* Beet

Blowfly (*Calliphora erythrocephala*), purification and properties of *o*-diphenol oxidase from third-instar larvae of (Pau, R. N. & Eagles, P. A. M.) 707–712

Bone, ox, mechanism of the stabilization of the reducible intermediate cross-links of collagens from, and other tissues (Robins, S. P. & Bailey, A. J.) 381–385

Brain, pig, regulatory effects of fatty acyl-coenzyme A on the activity of phosphate-activated glutaminase from, and kidney *in vitro* (Kvamme, E. & Torgner, I. Aa.) 83–91

Brain, rat, comparison of the activities of phosphatidylinositol phosphodiesterase from, towards phosphatidylinositol in dispersions and as monolayers at the air/water interface (Quinn, P. J. & Barenholz, Y.) 199–208

Brain, rat, discrimination of the antigenic site and the thiol-blocking-inhibitor-sensitive site of isoenzymes of hexokinase of mitochondria and cytosol of, and other tissues (Ouchi, M. & Ishibashi, S.) 481–483

Brain, rat, reciprocal exclusion by L-3,4-dihydroxyphenylalanine and L-tyrosine of their incorporation as single units into a soluble protein of (Rodríguez, J. A., Barra, H. S., Arce, C. A., Hallak, M. E. & Caputto, R. 115–121

Caffeate, kinetics of the hydroxylation of *p*-coumarate to, by beet-leaf phenolase (McIntrye, R. J. & Vaughan, P. F. T.) 447–461

Calcium ions, activation of sheep spermatozoal acrosin by, and their influence on the reaction of the enzyme with inhibitors (Brown, C. R., Andani, Z. & Hartree, E. F.) 133–146

Calcium ions, effect of, on the interaction of magnesium ions with teichoic acid from *Lactobacillus buchneri* (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 519–524

Calliphora erythrocephala, *see* Blowfly
Cartilage, laryngeal, pig, nature of the protein moieties of the proteoglycans from (Baxter, E. & Muir, H.). 657–668
Cartilage, nasal-septum, ox, isolation of a collagen-associated proteoglycan from, and its preferential interaction with α_2 chains of calf skin type I collagen (Lee-Own, V. & Anderson, J. C.) 57–63
Cartilage, nasal-septum, ox, nature of the protein moieties of the proteoglycans from (Baxter, E. & Muir, H.) 657–668
Cations, activation of sheep spermatozoal acrosin by, and their influence on the reaction of the enzyme with inhibitors (Brown, C. R., Andani, Z. & Hartree, E. F.) 133–146
Cell wall, *Lactobacillus buchneri* N.C.I.B. 8017, interaction of magnesium ions with teichoic acid of (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 519–524
Cell wall, *Pseudomonas aeruginosa* N.C.T.C. 8505, identification of 2-amino-2-deoxygalacturonic acid as a component of the lipopolysaccharide from (Wilkinson, S. G. & Welbourn, A. P.) 783–784
Cell wall, *Staphylococcus aureus* H mutant, bacteriophage-resistant, linkage between teichoic acid and peptidoglycan in (Heckels, J. E., Archibald, A. R. & Baddiley, J.) 637–647
Cell walls, detection of acetyl groups in preparations of, from grasses and other higher plants (Bacon, J. S. D., Gordon, A. H., Morris, E. J. & Farmer, V. C.) 485–487
Cells, epithelial, mammalian, photochemical binding of anthracene to deoxyribonucleic acid of, in tissue culture (Blackburn, G. M. & Taussig, P. E.) 289–291
Cells, plasmacytoma, MPC-11, mouse, determination of polyadenylate-rich ribonucleic acid in the nucleus and cytoplasm of (Abraham, K. A. & Eikhom, T. S.) 669–674
Cephalosporin C, reaction of, with pig insulin (Corran, P. H. & Waley, S. G.) 357–364
Chick embryo, analysis by two-dimensional gel electrophoresis of the proteins of ribosome crystals from (Dondi, P. G.) 475–476
Chicken, behaviour of mixed monolayers of fusogenic lipids and membrane phospholipids of erythrocytes from (Maggio, B. & Lucy, J. A.) 597–608
Chicken egg, electron-paramagnetic-resonance spectroscopy of iron-binding fragments of ovotransferrins from (Butterworth, R. M., Gibson, J. F. & Williams, J.) 559–563
Chicken, preparation and properties of iron-binding fragments from the *C*-terminal region of ovotransferrin from (Williams, J.) 237–244
Chloride ions, effects of, and other ions on the activity of monomeric (lobster) and dimeric (sea-cucumber) arginine kinases (Anosike, E. O. & Watts, D. C.) 387–395
Chlorophyll, organization of, in protein–chlorophyll complexes from pea leaves (Scott, B. & Gregory, R. P. F.) 341–347
Chloroplasts, pea-leaf, organization of chlorophyll in protein–chlorophyll complexes from (Scott, B. & Gregory, R. P. F.) 341–347
Cholesterol, evidence for a decarboxylation mechanism not involving a Schiff base intermediate in the conversion of 4,4-dimethylcholest-7-enol into 4α-methylcholest-7-enol during the biosynthesis of, by rat liver microsomal fraction (Wilton, D. C. & Akhtar, M.) 233–235
Chromatium sp., X-ray photoelectron spectra of high-potential iron–sulphur protein from, and of other iron–sulphur proteins (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474
Chromous ions, reduction by, of horse heart cytochrome *c* modified by reaction with 2-hydroxy-5-nitrobenzyl bromide (Brittain, T. & Greenwood, C.) 179–185
trans-Cinnamate, kinetics of the formation of, from L-phenylalanine catalysed by *Rhizoctonia solani* L-phenylalanine ammonia-lyase (Kalghatgi, K. K. & Subba Rao, P. V.) 65–72
Clearing-factor lipase, *see* Lipase, clearing-factor
Clostridium pasteurianum, X-ray photoelectron spectra of ferredoxin from, and of other iron–sulphur proteins (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474
Collagen, type I, skin, calf, isolation of a collagen-associated proteoglycan from ox nasal-septum cartilage and its preferential interaction with (Lee-Own, V. & Anderson, J. C.) 57–63
Collagens, ox, various, mechanism of the stabilization of the reducible intermediate cross-links of (Robins, S. P. & Bailey, A. J.) 381–385
Comamonas terrigena, occurrence of secondary alkylsulphatases in a strain of (Fitzgerald, J. W.) 477–480
Complement, guinea-pig, effect of reduction and fragmentation of rabbit immunoglobulin G on the fixation of component C1 of, by immune complexes (Press, E. M.) 285–288
Computer, method for the analysis by, of simple spectrophotometric changes in particulate systems exemplified by the effect of temperature on the 695nm absorption band of immobilized horse heart cytochrome *c* (Moore, T. A. & Greenwood, C.) 169–177
Connective tissues, mammalian, various, comparative studies of the cross-linked regions of elastins from (Gerber, G. E. & Anwar, R. A.) 685–695
p-Coumarate, kinetics of the formation of, from L-tyrosine catalysed by *Rhizoctonia solani* L-phenylalanine ammonia-lyase (Kalghatgi, K. K. & Subba Rao, P. V.) 65–72
p-Coumarate, kinetics of the hydroxylation of, to caffeate by beet-leaf phenolase (McIntyre, R. J. & Vaughan, P. F. T.) 447–461
Crithidia oncopelti, physicochemical properties of *Euglena gracilis* cytochrome *c*-558 and of cytochrome *c*-557 from (Pettigrew, G. W., Aviram, I. & Schejter, A.) 155–167
Cucumis melo, *see* Melon
Cyclic nucleotide phosphodiesterases, three, resolution, partial purification and properteis of, from potato tuber (Ashton, A. R. & Polya, G. M.) 329–339
Cytochrome *c*-557, *Crithidia oncopelti*, physicochemical properties of *Euglena gracilis* cytochrome *c*-558 and (Pettigrew, G. W., Aviram, I. & Schejter, A.) 155–167
Cytocrome *c*-553 (cytochrome *f*), *Alaria esculenta*, amino acid sequence of (Laycock, M. V.) 271–279

Cytochrome *c*-558, *Euglena gracilis*, physicochemical properties of *Crithidia oncopelti* cytochrome *c*-557 and (Pettigrew, G. W., Aviram, I. & Schejter, A.) 155–167
Cytochrome *c*, heart, horse, immobilized, method for the investigation of the effect of temperature on the 695 nm absorption band of (Moore, T. A. & Greenwood, C.) 169–177
Cytochrome *c*, heart, horse, kinetic studies on, *N*-formylated at its tryptophan residue (Brittain, T. & Greenwood, C.) 713–717
Cytochrome *c*, heart, horse, kinetic studies on, modified by reaction with 2-hydroxy-5-nitrobenzyl bromide (Brittain, T. & Greenwood, C.) 179–185
Cytochrome *f*, *Alaria esculenta*, amino acid sequence of (Laycock, M. V.) 271–279
Cytosol, brain, rat, discrimination of the antigenic site and the thiol-blocking-inhibitor-sensitive site of isoenzymes of hexokinase of, and mitochondria and of other tissues (Ouchi, M. & Ishibashi, S.) 481–483
Cytosol, liver, rat, reduction of disulphide bonds in proteins and protein mixed disulphides by a thioltransferase in (Mannervik, B. & Axelsson, K.) 785–788
Cytosol, plasmacytoma, MPC-11, mouse, determination of polyadenylate-rich ribonucleic acid in, and nuclei (Abraham, K. A. & Eikhom, T. S.) 669–674

Datura innoxia roots, oxidation of ethyl formylphenylacetate by oxidase–peroxidase enzymes from (Kalyanaraman, V. S., Mahadevan, S. & Kumar, S. A.) 565–576
Datura innoxia roots, oxidation of reduced nicotinamide–adenine dinucleotide in the presence of ethyl formylphenylacetate by oxidase–peroxidase enzymes from (Kalyanaraman, V. S., Kumar, S. A. & Mahadevan, S.) 577–584
Dehydrodihydroxylysinonorleucine, mechanism of the stabilization of the reducible intermediate cross-links dehydrohydroxylysinonorleucine and, in various ox collagens (Robins, S. P. & Bailey, A. J.) 381–385
Dehydrogenases, nicotinamide–adenine dinucleotide-dependent, sequential, stereospecificity of, in relation to the evolution of metabolic sequences (do Nascimento, K. H. & Davies, D. D.) 553–557
Dehydrogenases, use of a double-beam rapid-scanning stopped-flow spectrophotometer to measure rapid changes in the spectrum of reduced nicotinamide–adenine dinucleotide during reactions catalysed by (Hollaway, M. R. & White, H. A.) 221–231
Dehydrohydroxylysinonorleucine, mechanism of the stabilization of the reducible intermediate cross-links dehydrodihydroxylysinonorleucine and, in various ox collagens (Robins, S. P. & Bailey, A. J.) 381–385
Deoxyribonucleic acid, photochemical binding of anthracene to, of monkey kidney epithelial cells and human skin epithelial cells in tissue culture (Blackburn, G. M. & Taussig, P. E.) 289–291
Deoxyribonucleic acid, relationship between *Halobacterium cutirubrum* N.R.C. 34001, deoxyribonucleic acid-depenedent ribonucleic acid polymerase bound to, and the low-molecular-weight soluble form of the enzyme (Fitt, P. S., Peterkin, P. I. & Barua, N. N.) 719–724
Deoxyribonucleic acid, satellite, melon, complexity of (Sinclair, J., Wells, R., Duemling, B. & Ingle, J.) 31–38
Desmosine cross-links, comparative studies of, of elastins from ox ligamentum nuchae and ox, pig and human aorta (Gerber, G. E. & Anwar, R. A.) 685–695
Diet, induction of metallothionein in rat liver by the injection of zinc ions and by the restriction of the amount of (Bremner, I. & Davies, N. T.) 733–738
L-3,4-Dihydroxyphenylalanine (L-dopa), reciprocal exclusion by L-tyrosine and, of their incorporation as single units into a rat brain soluble protein (Rodríguez, J. A., Barra, H. S., Arce, C. A., Hallak, M. E. & Caputto, R.) 115–121
4,4-Dimethylcholest-7-enol, evidence for a decarboxylation mechanism not involving a Schiff base intermediate in the conversion of, into 4α-methylcholest-7-enol by rat liver microsomal fraction (Wilton, D. C. & Akhtar, M.) 233–235
Dinucleotides, calculations of the conformational energy of adenosine 5′-monophosphate, nicotinamide mononucleotide and adenosine 3′-monophosphate moieties in (Thornton, J. M. & Bayley, P. M.) 585–596
o-Diphenol oxidase, purification and properties of, from third-instar larvae of the blowfly *Calliphora erythrocephala* (Pau, R. N. & Eagles, P. A. M.) 707–712
Disulphide bonds, effect of reduction of, or rabbit immunoglobulin G on the fixation of component C1 of guinea pig complement by immune complexes (Press, E. M.) 285–288
Disulphide bonds, evidence against a proposed interaction of thionitrobenzoate with, of proteins (Weitzman, P. D. J.) 281–283
Disulphide bonds, reduction of, in proteins and protein mixed disulphides by a thioltransferase in rat liver cytosol (Mannervik, B. & Axelsson, K.) 785–788
Disulphides, effects of various amine oxides and, on the activity of ox erythrocyte acetylcholinesterase (Dawson, R. M.) 293–295
L-Dopa (L-3,4-dihydroxyphenylalanine), reciprocal exclusion by L-tyrosine and, of their incorporation as single units into a rat brain soluble protein (Rodríguez, J. A., Barra, H. S., Arce, C. A., Hallak, M. E. & Caputto, R.) 115–121

Egg, chicken, electron-paramagnetic-resonance spectroscopy of iron-binding fragments of ovotransferrin from (Butterworth, R. M., Gibson, J. F. & Williams, J.) 559–563
Egg-white, chicken, reaction of benzylpenicillin with lysozyme from (Corran, P. H. & Waley, S. G.) 357–364
Egg-white, chicken, stopped-flow fluorescence studies on the binding of the β-(1→4)-linked trimer of *N*-acetyl-D-glucosamine to lysozyme from (Halford, S. E.) 411–422
Elastins, comparative studies of the cross-linked regions of, from ox ligamentum nuchae and ox, pig and human aorta (Gerber, G. E. & Anwar, R. A.) 685–695
Embryo, chick, analysis by two-dimensional gel electrophoresis of the proteins of ribosome crystals from (Dondi, P. G.) 475–476

Enzymes, comparison of seven methods for fitting the Michaelis–Menten equation for reactions catalysed by (Atkins, G. L. & Nimmo, I. A.) 775–777
Enzymes, sigmoid curves, non-linear double-reciprocal plots and allosterism in reactions catalysed by (Bardsley, W. G. & Childs, R. E.) 313–328
Enzymes, use of the direct linear plot for determining initial velocities of reactions catalysed by (Cornish-Bowden, A.) 305–312
Epidermis, ox, extraction and characterization of α-keratin from (Steinert, P. M.) 39–48
Erythrocruorin, snail, physicochemical properties of (Wood, E. J. & Mosby, L. J.) 437–445
Erythrocytes, chicken, behaviour of mixed monolayers of fusogenic lipids and phospholipids of (Maggio, N. & Lucy, J. A.) 597–608
Erythrocytes, ox, effects of some amine oxides and disulphide compounds on the activity of acetylcholinesterase from (Dawson, R. M.) 293–295
Escherichia coli, interaction of thiol-containing analogues of puromycin with *N*-acetylphenylalanyl-transfer ribonucleic acid on ribosomes from rat liver and, to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220
Ethyl formylphenylacetate, oxidation of, by oxidase–peroxidase enzymes from *Datura innoxia* roots (Kalyanaraman, V. S., Mahadevan, S. & Kumar, S. A.) 565–576
Ethyl formylphenylacetate, oxidation of reduced nicotinamide–adenine dinucleotide in the presence of, by oxidase–peroxidase enzymes from *Datura innoxia* roots (Kalyanaraman, V. S., Kumar, S. A. & Mahadevan, S.) 577–584
Euglena gracilis, physiochemical properties of *Crithidia oncopelti* cytochrome *c*-557 and of cytochrome *c*-558 from (Pettigrew, G. W., Aviram, I. & Schejter, A.) 155–167

Fatty acyl-coenzyme A, regulatory effects of, on the activity of pig brain and kidney phosphate-activated glutaminase *in vitro* (Kvamme, E. & Torgner, I.Aa.) 83–91
Ferredoxin, *Clostridium pasteurianum*, X-ray photoelectron spectra of, and other iron–sulphur proteins (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474
Ferredoxin, spinach, X-ray photoelectron spectra of, and other iron–sulphur proteins (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474
Ferricytochrome *c*, heart, horse, immobilized, method for the investigation of the effect of temperature on the 695nm absorption band of (Moore, T. A. & Greenwood, C.) 169–177
Flour, wheat, structural studies on three proteins from, that lack phenylalanine and histidine residues (Redman, D. G.) 725–732
Follicle-stimulating-hormone receptor, properties of, of ox testis plasma membrane (Cheng, K.-W.) 123–132
N-Formylation, effects of, of the tryptophan residue of horse heart cytochrome *c* on the kinetic properties of the protein (Brittain, T. & Greenwood, C.) 713–717
Formylphenylacetic acid ethyl ester, oxidation of, by oxidase–peroxidase enzymes from *Datura innoxia* roots (Kalyanaraman, V. S., Mahadevan, S. & Kumar, S. A.) 565–576
Formylphenylacetic acid ethyl ester, oxidation of reduced nicotinamide–adenine dinucleotide in the presence of, by oxidase–peroxidase enzymes from *Datura innoxia* roots (Kalyanaraman, V. S., Kumar, S. A. & Mahadevan, S.) 577–584

Gadolinium ions, binding of, to non-immune rabbit immunoglobulin G and its fragments (Dower, S. K., Dwek, R. A., McLaughlin, A. C., Mole, L. E., Press, E. M. & Sunderland, C. A.) 73–82
Glutamate dehydrogenase, liver, ox, role of lysine residue 126 in the activity of, in the light of studies of the inhibition and inactivation of the enzyme by pyridoxal 5′-phosphate (Chen, S.-S. & Engel, P. C.) 619–626
Glutamate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-specific, *Neurospora crassa*, amino acid sequence of the blocked *N*-terminal peptide of (Morris, H. R. & Dell, A.) 754–755
Glutamate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-specific *Neurospora crassa*, amino acid sequences of peptides obtained by the digestion of, with a staphylococcal proteinase (Wootton, J. C., Baron, A. J. & Fincham, J. R. S.) 749–754
Glutamate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-specific, *Neurospora crassa*, amino acid sequences of peptides obtained by the digestion of, with pepsin and chymotrypsin, and the complete amino acid sequence of the enzyme (Holder, A. A., Wootton, J. C., Baron, A. J., Chambers, G. K. & Fincham, J. R. S.) 757–773
Glutamate dehydrogenase, nicotinamide–adenine dinucleotide phosphate-specific, *Neurospora crassa*, amino acid sequences of peptides obtained by the digestion of, with trypsin (Wootton, J. C., Taylor, J. G., Jackson, A. A., Chambers, G. K. & Fincham, J. R. S.) 739–748
Glutamate, glutamine and asparagine as nitrogen donors for the reductant-dependent biosynthesis of, in pea roots (Miflin, B. J. & Lea, P. J.) 403–409
Glutaminase, phosphate-activated, brain and kidney, regulatory effects of fatty acyl-coenzyme A on the activity of, *in vitro* (Kvamme, E. & Torgner, I. Aa) 83–91
Glutamine, asparagine and, as nitrogen donors for the reductant-dependent biosynthesis of glutamate in pea roots (Miflin, B. J. & Lea, P. J.) 403–409
Glutathione, reduced, inhibition by, of the hydrolysis of glycerol 2-phosphate by pig kidney alkaline phosphatase (Ahlers, J.) 535–546
Glyceraldehyde 3-phosphate dehydrogenase, skeletal-muscle, rabbit, use of a double-beam rapid-scanning stopped-flow spectrophotometer to measure rapid changes in the spectrum of reduced nicotinamide–adenine dinucleotide during the reaction catalysed by (Hollaway, M. R. & White, H. A.) 221–231
Glycerol 2-phosphate, mechanism of the hydrolysis of, by pig kidney alkaline phosphatase (Ahlers, J.) 535–546

Glycerol phosphate residues, presence of, in the linkage between teichoic acid and peptidoglycan in the cell wall of a bacteriophage-resistant mutant of *Staphylococcus aureus* H (Heckels, J. E., Archibald, A. R. & Baddiley, J.) 637–647

β-Glycerophosphate, *see* Glycerol 2-phosphate

Glycolysis, identification in human parotid-gland saliva of a peptide particularly active in enhancing, by salivary micro-organisms (Holbrook, I. B. & Molan, P. C.) 489–492

Grasses, detection of acetyl groups in preparations of cell walls from, and other higher plants (Bacon, J. S. D., Gordon, A. H., Morris, E. J. & Farmer, V. C.) 485–487

Haemoglobin (erythrocruorin), snail, physicochemical properties of (Wood, E. J. & Mosby, L. J.) 437–445

Haemoglobin, rat, major, partial amino acid sequence of the α chain of (Chua, C. G., Carrell, R. W. & Howard, B. H.) 259–269

Haemoglobins, rat, identification of two structurally distinct α chains and functional behaviour of selected components of (Garrick, L. M., Sharma, V. S., McDonald, M. J. & Ranney, H. M.) 245–258

Halobacterium cutirubrum N.R.C. 34001, relationship between the deoxyribonucleic acid-bound and low-molecular-weight soluble forms of deoxyribonucleic acid-dependent ribonucleic acid polymerase from (Fitt, P. S., Peterkin, P. I. & Barua, N. N.) 719–724

Heart, horse, kinetic studies on cytochrome *c* from, *N*-formylated at its tryptophan residue (Brittain, T. & Greenwood, C.) 713–717

Heart, horse, kinetic studies on cytochrome *c* from, modified by reaction with 2-hydroxy-5-nitrobenzyl bromide (Brittain, T. & Greenwood, C.) 179–185

Heart, horse, method for the investigation of the effect of temperature on the 695nm absorption band of immobilized cytochrome *c* from (Moore, T. A. & Greenwood, C.) 169–177

Heart muscle, pig, use of digestion with *Armillaria mellea* proteinase having specificity for lysine residues in the establishment of the amino acid sequence of aspartate aminotransferase from (Doonan, S., Doonan, H. J., Hanford, R., Vernon, C. A., Walker, J. M., Airoldi, L. P. da S., Bossa, F., Barra, D., Carloni, M., Fasella, P. & Riva, F.) 497–506

Heart muscle, rabbit and ox, phosphorylation of troponin I from (Cole, H. A. & Perry, S. V.) 525–533

Heart, pig, purification and characterization of clearing-factor lipase from myocardium from (Ehnholm, C Kinnunen, P. K. J., Huttunen, J. K., Nikkilä, E. A. & Ohta, M.) 649–655

Hen, *see* Chicken

Heparin, solubilization and partial characterization of *N*- and *O*-sulphotransferases involved in the biosynthesis of, from mouse mastocytoma microsomal fraction (Jansson, L., Höök, M., Wasteson, Å, & Lindahl, U.) 49–55

Hexokinase, mitochondrial and cytosolic, discrimination of the antigenic site and the thiol-blocking-inhibitor-sensitive site of isoenzymes of, or rat brain and other tissues (Ouchi, M. & Ishabishi, S.) 481–483

Histidine residues, structural studies on three wheat-flour proteins that lack phenylalanine residues and (Redman, D. G.) 725–732

Holothuria forskali, *see* Sea-cucumber

Homarus vulgaris, *see* Lobster

Housefly (*Musca domestica*), properties of acetylcholinesterase from, resistant and susceptible to organophosphorus insecticides (Devonshire, A. L.) 463–469

Hyaluronidase, nature of the protein moieties of the proteoglycans of cartilage from pig larynx and ox nasal septum released by digestion with (Baxter, E. & Muir, H.) 657–668

2-Hydroxy-5 nitrobenzyl bromide, kinetic studies on horse heart cytochrome *c* modified by reaction with (Brittain, T. & Greenwood, C.) 179–185

Immunoglobulin G, rabbit, effect of reduction and fragmentation of, on the fixation of component Cl of guinea-pig complement by immune complexes (Press, E. M.) 285–288

Immunoglobulin G, rabbit, non-immune, binding of lanthanide ions to, and its fragments (Dower, S. K., Dwek, R. A., McLaughlin, A. C., Mole, L. E., Press, E. M. & Sunderland, C. A.) 73–82

Insecticides, organophosphorus, properties of acetylcholinesterase from houseflies resistant and susceptible to (Devonshire, A. L.) 463–469

Insulin, pig, reaction of benzylpenicillin with (Corran, P. H. & Waley, S. G.) 357–364

Iron, electron-paramagnetic-resonance spectroscopy of fragments of chicken ovotransferrin binding (Butterworth, R. M., Gibson, J. F. & Williams, J.) 559–563

Iron, preparation and properties of fragments binding, from the *C*-terminal region of chicken ovotransferrin (Williams, J.) 237–244

Iron–sulphur protein, high-potential, *Chromatium* sp., X-ray photoelectron spectra of, and other iron–sulphur proteins (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474

Iron–sulphur proteins, X-ray photoelectron spectra of (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474

Isoniazid (isonicotinic acid hydrazide), purification and properties of peroxidase from *Mycobacterium tuberculosis* H37Rv and its possible role in the mechanism of action of, on the organism (Gayathri Devi, B., Shaila, M. S., Ramakrishnan, T. & Gopinathan, K. P.) 187–197

Isonicotinic acid hydrazide (isoniazid), purification and properties of peroxidase from *Mycobacterium tuberculosis* H37Rv and its possible role in the mechanism of action of, on the organism (Gayathri Devi, B., Shaila, M. S., Ramakrishnan, T. & Gopinathan, K. P.) 187–197

α-Keratin, extraction and characterization of, from ox epidermis (Steinert, P. M.) 39–48

Kidney epithelial cells, monkey, photochemical binding of anthracene to deoxyribonucleic acid of human skin epithelial cells and, in tissue culture (Blackburn, G. M. & Taussig, P. E.) 289–291

Kidney, pig, mechanism of the hydrolysis of glycerol 2-phosphate by alkaline phosphatase from (Ahlers, J.) 535–546
Kidney, pig, regulatory effects of fatty acyl-coenzyme A on the activity of phosphate-activated glutaminase from, and brain *in vitro* (Kvamme, E. & Torgner, I. Aa.) 83–91
Kinase II, *see* Angiotensin-converting enzyme

β-Lactamase I, *Bacillus cereus* 569/H, dependence on pH of the hydrolysis of benzylpenicillin and ampicillin by (Waley, S. G.) 547–551
β-Lactamase, *Staphylococcus aureus* PC1, isolation of a specific covalent complex of, with quinacillin (Virden, R., Bristow, A. F. & Pain, R. H.) 397–401
Lactate dehydrogenase, muscle, pig, inactivation of the M_4 isoenzyme of, by reaction of pyridoxal 5′-phosphate with an essential lysine residue (Chen, S.-S. & Engel, P. C.) 107–113
Lactobacillus buchneri N.C.I.B. 8007, interaction of magnesium ions with teichoic acid from (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 519–524
Lanthanide ions, binding of, to non-immune rabbit immunoglobulin G and its fragments (Dower, S. K., Dwek, R. A., McLaughlin, A. C., Mole, L. E., Press, E. M. & Sunderland, C. A.) 73–82
Larynx, pig, nature of the protein moieties of the proteoglycans of cartilage from ox nasal septum and (Baxter, E. & Muir, H.) 657–668
Leaves, beet, kinetics of the hydroxylation of *p*-coumarate to caffeate by phenolase from (McIntrye, R. J. & Vaughan, P. F. T.) 447–461
Leaves, pea, organization of chlorophyll in protein–chlorophyll complexes from (Scott, B. & Gregory, R. P. F.) 341–347
Lecithin, *see* Phosphatidylcholine
Ligamentum nuchae, ox, comparative studies of the cross-linked regions of elastins from, and ox, pig and human aorta (Gerber, G. E. & Anwar, R. A.) 685–695
Linoleoyl-coenzyme A, regulatory effects of, and other fatty acyl-coenzyme A species on the activity of pig brain and kidney phosphate-activated glutaminase *in vitro* (Kvamme, E. & Torgner, I. Aa.) 83–91
Lipase, clearing-factor, purification and characterization of, from pig myocardium (Ehnholm, C., Kinnunen, P. K. J., Huttenen, J. K., Nikkilä, E. A. & Ohta, M.) 649–655
Lipids, fusogenic, behaviour of mixed monolayers of phospholipids and (Maggio, B. & Lucy, J. A.) 597–608
Lipopolysaccharide, *Pseudomonas aeruginosa* N.C.T.C. 1999, characterization of polysaccharide fractions obtained by mild acid hydrolysis of (Drewry, D. T., Symes, K. C., Gray, G. W. & Wilkinson, S. G.) 93–106
Lipopolysaccharide, *Pseudomonas aeruginosa* N.C.T.C. 8505, identification of 2-amino-2-deoxygalacturonic acid as a component of (Wilkinson, S. G. & Welbourn, A. P.) 783–784
Lipoprotein lipase, *see* Lipase, clearing-factor
Lipoproteins, apo-, serum, guinea-pig, distribution and partial characterization of (Chapman, M. J., Mills, G. L. & Ledford, J. H.) 423–436
Liver, guinea-pig, stereospecificity of nicotinamide–adenine dinucleotide-dependent xylitol dehydrogenase from, and of other dehydrogenases (do Nascimento, K. H. & Davies, D. D.) 553–557
Liver, horse, inactivation of alcohol dehydrogenase from, by reaction of pyridoxal 5′-phosphate with the essential lysine residue (Chen, S.-S. & Engel, P. C.) 627–635
Liver, horse, use of a double-beam rapid-scanning stopped-flow spectrophotometer to measure rapid changes in the spectrum of reduced nicotinamide–adenine dinucleotide during the reaction catalysed by alcohol dehydrogenase from (Hollaway, M. R. & White, H. A.) 221–231
Liver, ox, role of lysine residue 126 in the activity of glutamate dehydrogenase from, in the light of studies of the inhibition and inactivation of the enzyme by pyridoxal 5′-phosphate (Chen, S.-S. & Engel, P. C.) 619–626
Liver, rat, evidence for a decarboxylation mechanism not involving a Schiff base intermediate in the conversion of 4,4-dimethylcholest-7-enol into 4α-methylcholest-7-enol by microsomal fraction from (Wilton, D.C. & Akhtar, M.) 233–235
Liver, rat, induction of metallothionein in, by the injection of zinc ions and by the restriction of food intake (Bremner, I. & Davies, N. T.) 733–738
Liver, rat, interaction of thiol-containing analogues of puromycin with *N*-acetylphenylanyl-transfer ribonucleic acid on ribosomes from *Escherichia coli* and, to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220
Liver, rat, reduction of disulphide bonds in proteins and protein mixed disulphides by a thioltransferase in cytosol from (Mannervik, B. & Axelsson, K.) 785–788
Liver, sheep, purification of 2-oxo aldehyde dehydrogenase from, and its dependence on unusual amines (Dunkerton, J. & James, S. P.) 609–617
Lobster (*Homarus vulgaris*), effects of anions on the activity of arginine kinases from sea-cucumber (*Holothuria forskali*) and (Anosike, E. O. & Watts, D. C.) 387–395
Lung, pig, sensitive radiochemical method for assay of the activity of angiotensin-converting enzyme (kinase II) from (Chiu, A. T., Ryan, J. W., Ryan, U. S. & Dorer, F. E.) 297–300
Lysine residue, essential, inactivation of horse liver alcohol dehydrogenase by reaction of pyridoxal 5′-phosphate with (Chen, S.-S. & Engel, P. C.) 627–635
Lysine residue, essential, inactivation of pig muscle lactate dehydrogenase M_4 isoenzyme by reaction of pyridoxal 5′-phosphate with (Chen, S.-S. & Engel, P. C.) 107–113
Lysine residue 126, role of, in the activity of ox liver glutamate dehydrogenase in the light of studies of the inhibition and inactivation of the enzyme by pyridoxal 5′-phosphate (Chen, S.-S. & Engel, P. C.) 619–626
Lysine residues, use of digestion with *Armillaria mellea* proteinase having specificity for, in the establishment of the amino acid sequence of pig heart-muscle aspartate aminotransferase (Doonan, S., Doonan, H. J., Hanford, R., Vernon, C. A., Walker, J. M., Airoldi, L. P. da S., Bossa, F., Barra, D., Carloni, M., Fasella, P. & Riva, F.) 497–506

Lysozyme, hen's-egg-white, reaction of benzylpenicillin with (Corran, P. H. & Waley, S. G.) 357–364
Lysozyme, hen's-egg-white, stopped-flow fluorescence studies on the binding of the β-(1→4)-linked trimer of *N*-acetyl-D-glucosamine to (Halford, S. E.) 411–422

Magnesium ions, activation by, of the hydrolysis of glycerol 2-phosphate by pig kidney alkaline phosphatase (Ahlers, J.) 535–546
Magnesium ions, interaction of, with teichoic acid from *Lactobacillus buchneri* N.C.I.B. 8007 (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 519–524
Malate dehydrogenase, stereospecificity of, and of nicotinamide–adenine dinucleotide-dependent dehydrogenases associated with the metabolism of *meso*-tartrate in *Pseudomonas putida* A.T.C.C. 17642 (do Nascimento, K. H. & Davies, D. D.) 553–557
'Malic' enzyme (EC 1.1.1.40), potato-tuber, unidirectional inhibition and activation by *meso*-tartrate of (do Nascimento, K. H., Davies, D. D. & Patil, K. D.) 349–355
Mannitol 1-phosphate dehydrogenase, nicotinamide–adenine dinucleotide-dependent, *Aerobacter aerogenes* N.C.I.B. 8017, stereospecificity of, and sorbitol 6-phosphate dehydrogenase (do Nascimento, K. H. & Davies, D. D.) 553–557
Mastocytoma, mouse, solubilization and partial characterization of *N*- and *O*-sulphotransferases involved in the biosynthesis of heparin from microsomal fraction of (Jansson, L., Höök, M., Wasteson, Å. & Lindahl, U.) 49–55
Melon (*Cucumis melo*), complexity of satellite deoxyribonucleic acid in (Sinclair, J., Wells, R., Deumling, B. & Ingle, J.) 31–38
Membrane, erythrocyte, chicken, behaviour of mixed monolayers of fusogenic lipids and phospholipids of (Maggio, B. & Lucy, J. A.) 597–608
Membrane, plasma, testis, ox, properties of follicle-stimulating-hormone receptor of (Cheng, K.-W.) 123–132
Messenger ribonucleic acid, *see* Ribonucleic acid, messenger
Metallothionein, induction of, in rat liver by the injection of zinc ions and by the restriction of food intake (Bremner, I. & Davies, N. T.) 733–738
4α-Methylcholest-7-enol, evidence for a decarboxylation mechanism not involving a Schiff base intermediate in the conversion of 4,4-dimethylcholest-7-enol into, by rat liver microsomal fraction (Wilton, D. C. & Akhtar, M.) 233–235
4-Methylumbelliferyl *p*-guanidinobenzoate, determination of the operational molarity of sheep spermatozoal acrosin by fluorimetric titration with (Brown, C. R., Andani Z. & Hartree, E. F.) 147–154
Michaelis–Menten equation, comparison of seven methods for fitting, for reactions catalysed by enzymes (Atkins, G. L. & Nimmo, I. A.) 775–777
Microsomal fraction, liver, rat, evidence for a decarboxylation mechanism not involving a Schiff base intermediate in the conversion of 4,4-dimethylcholest-7-enol into 4α-methylcholest-7-enol by (Wilton, D. C. & Akhtar, M.) 233–235
Microsomal fraction, mastocytoma, mouse, solubilization and partial characterization of *N*- and *O*-sulphotransferases involved in the biosynthesis of heparin from Jansson, L., Höök, M., Wasteson, Å. & Lindahl, U.) 49–55
Microtubules, brain, rat, reciprocal exclusion by 3,4-dihydroxyphenylalanine and L-tyrosine of their incorporation as single units into a soluble protein tentatively identified as a component of (Rodríguez, J. A., Barra, H. S., Arce, C. A., Hallak, M. E. & Caputto, R.) 115–121
Mitochondria, brain, rat, discrimination of the antigenic site and the thiol-blocking-inhibitor-sensitive site of isoenzymes of hexokinase of, and cytosol and of other tissues (Ouchi, M. & Ishibashi, S.) 481–483
Mitochondria, yeast, messenger activity of ribonucleic acid from (Eggitt, M. J. & Scragg, A. H.) 507–512
Musca domestica, *see* Housefly
Muscle, heart, pig, use of digestion with *Armillaria mellea* proteinase having specificity for lysine residues in the establishment of the amino acid sequence of aspartate aminotransferase from (Doonan, S., Doonan, H. J., Hanford, R., Vernon, C. A., Walker, J. M., Airoldi, L. P. da S., Bossa, F., Barra, D., Carloni, M., Fasella, P. & Riva, F.) 497–506
Muscle, heart, rabbit and ox, phosphorylation of troponin I from (Cole, H. A. & Perry, S. V.) 525–533
Muscle, pig, inactivation of lactate dehydrogenase M_4 isoenzyme from, by reaction of pyridoxal 5′-phosphate with an essential lysine residue (Chen, S.-S. & Engel, P. C.) 107–113
Muscle, skeletal, rabbit, amino acid sequence of troponin I from (Wilkinson, J. M. & Grand, R. J. A.) 493–496
Muscle, skeletal, rabbit, phosphorylation of troponin I from rabbit and ox heart muscle and from (Cole, H. A. & Perry, S. V.) 525–533
Muscle, skeletal, rabbit, use of a double-beam rapid-scanning stopped-flow spectrophotometer to measure rapid changes in the spectrum of reduced nicotinamide–adenine dinucleotide during the reaction catalysed by glyceraldehyde 3-phosphate dehydrogenase from (Hollaway, M. R. & White, H. A.) 221–231
Muscle, skeletal, rabbit, use of affinity chromatography for assessment of the interactions of actin and myosin from (Bottomley, R. C. & Trayer, I. P.) 365–379
Mycobacterium tuberculosis H37Rv, purification and properties of peroxidase from, and its possible role in the mechanism of action of isonicotinic acid hydrazide on the organism (Gayathri Devi, B., Shaila, M. S., Ramakrishnan, T. & Gopinathan, K. P.) 187–197
Myocardium, pig, purification and characterization of clearing-factor lipase from (Ehnholm, C., Kinnunen, P. K. J., Huttunen, J. K., Nikkilä, E. A. & Ohta, M.) 649–655
Myosin, skeletal-muscle, rabbit, use of affinity chromatography for assessment of the interactions of, and actin (Bottomley, R. C. & Trayer, I. P.) 365–379

Nasal septum, ox, isolation of a collagen-associated proteoglycan from cartilage of, and its preferential interaction with α_2 chains of calf skin type I collagen (Lee-Own, V. & Anderson, J. C.) 57–63

Nasal septum, ox, nature of the protein moieties of the proteoglycans of cartilage from pig larynx and (Baxter, E. & Muir, H.) 657–668

Neurospora crassa, amino acid sequence of the blocked *N*-terminal peptide of nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase from (Morris, H. R. & Dell, A.) 754–755

Neurospora crassa, amino acid sequences of peptides obtained by the digestion of nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase from, with pepsin and chymotrypsin, and the complete amino acid sequence of the enzyme (Holder, A. A., Wotton, J. C., Baron, A. J., Chambers, G. K. & Fincham, J. R. S.) 757–773

Neurospora crassa, amino acid sequences of peptides obtained by the digestion of nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase, with a staphylococcal proteinase (Wootton, J. C., Baron, A. J. & Fincham, J. R. S.) 749–754

Neurospora crassa, amino acid sequences of peptides obtained by the digestion of nicotinamide–adenine dinucleotide phosphate-specific glutamate dehydrogenase from, with trypsin (Wootton, J. C., Taylor, J. G., Jackson, A. A., Chambers, G. K. & Fincham, J. R. S.) 739–748

Nicotinamide–adenine dinucleotide (oxidized) pyrophosphatase, resolution, partial purification and characterization of three cyclic nucleotide phosphodiesterases possessing activity of, from potato tuber (Ashton, A. R. & Polya, G. M.) 329–339

Nicotinamide–adenine dinucleotide, reduced oxidation of, in the presence of ethyl formylphenylacetate by oxidase–peroxidase enzymes from *Datura innoxia* roots (Kalyanaraman V. S., Kumar, S. A. & Mahadevan, S.) 577–584

Nicotinamide–adenine dinucleotide, reduced, use of a double-beam rapid-scanning stopped-flow spectrophotometer to measure rapid changes in the spectrum of, during reactions catalysed by dehydrogenases (Hollaway, M. R. & White, H. A.) 221–231

Nicotinamide mononucleotide, calculations of the conformational energy of, and other mononucleotide moieties in dinucleotides (Thornton, J. M. & Bayley, P. M.) 585–596

Nitrate ions, effects of, and other anions on the activity of monomeric (lobster) and dimeric (sea-cucumber) arginine kinases (Anosike, E. O. & Watts, D. C.) 387–395

Nitrogen, glutamine and asparagine as donors of, for the reductant-dependent biosynthesis of glutamate in pea roots (Miflin, B. J. & Lea, P. J.) 403–409

Nuclei, plasmacytoma, MPC-11, mouse, determination of polyadenylate-rich ribonucleic acid in, and cytoplasm (Abraham, K. A. & Eikhom, T. S.) 669–674

O antigen, characterization of polysaccharide fractions obtained by mild acid hydrolysis of *Pseudomonas aeruginosa* N.C.T.C. 1999 lipopolysaccharide active as (Drewry, D. T., Symes, K. C., Gray, G. W. & Wilkinson, S. G.) 93–106

Oleoyl-coenzyme A, regulatory effects of, and other fatty acyl-coenzyme A species on the activity of pig brain and kidney phosphate-activated glutaminase *in vitro* (Kvamme, E. & Torgner, I. Aa.) 83–91

Oocytes, *Xenopus laevis*, messenger activity of yeast mitochondrial ribonucleic acid injected into (Eggitt, M. J. & Scragg, A. H.) 507–512

Organophosphorus insecticides, properties of acetylcholinesterase from houseflies resistant and susceptible to (Devonshire, A. L.) 463–469

Oriza sativa, *see* Rice

Ovotransferrin, chicken, electron-paramagnetic-resonance spectroscopy of iron-binding fragments of (Butterworth, R. M., Gibson, J. F. & Williams, J.) 559–563

Ovotransferrin, chicken, preparation and properties of iron-binding fragments from the *C*-terminal region of (Williams, J.) 237–244

Oxidase, oxidation of ethyl formylphenylacetate by an enzyme system possessing activities of peroxidase and, from *Datura innoxia* roots (Kalyanaraman, V. S., Mahadevan, S. & Kumar, S. A.) 565–576

Oxidase, oxidation of reduced nicotinamide–adenine dinucleotide in the presence of ethyl formylphenylacetate by an enzyme system possessing activities of peroxidase and, from *Datura innoxia* roots (Kalyanaraman, V. S., Kumar, S. A. & Mahadevan, S.) 577–584

Oxidoreductases, nicotinamide–adenine dinucleotide-dependent, sequential, stereospecificity of, in relation to the evolution of metabolic sequences (do Nascimento, K. H. & Davies, D. D.) 553–557

2-Oxo aldehyde dehydrogenase, purification of, from sheep liver and its dependence on unusual amines (Dunkerton, J. & James, S. P.) 609–617

Oxygen, binding of, to snail erythrocruorin (Wood, E. J. & Mosby, L. J.) 437–445

Palmitoyl-coenzyme A, regulatory effects of, and other fatty acyl-coenzyme A species on the activity of pig brain and kidney phosphate-activated glutaminase *in vitro* (Kvamme, E. & Torgner, I. Aa.) 83–91

Parotid gland, human, identification in saliva from, of a peptide particularly active in enhancing the glycolytic activity of salivary micro-organisms (Holbrook, I. B. & Molan, P. C.) 489–492

Pea (*Pisum sativum*) leaves, organization of chlorophyll in protein–chlorophyll complexes from (Scott, B. & Gregory, R. P. F.) 341–347

Pea (*Pisum sativum*) roots, glutamine and asparagine as nitrogen donors for the reductant-dependent biosynthesis of glutamate in (Miflin, B. J. & Lea, P. J.) 403–409

Penicillin, dependence on pH of the hydrolysis of derivatives of, by *Bacillus cereus* 569/H β-lactamase I (Waley, S. G.) 547–551

Penicillin, reaction of, with pig insulin and with hen's-egg-white lysozyme (Corran, P. H. & Waley, S. G.) 357–364

Penicillinase, *Bacillus cereus* 569/H, dependence on pH of the hydrolysis of benzylpenicillin and ampicillin by (Waley, S. G.) 547–551

Penicillinase, *Staphylococcus aureus* PC1, isolation of a specific covalent complex of, with quinacillin (Virden, R., Bristow, A. F. & Pain, R. H.) 397–401

Peptide, identification in human parotid-gland saliva of, particularly active in enhancing the glycolytic activity of salivary micro-organisms (Holbrook, I. B. & Molan, P. C.) 489–492

Peptides, nomenclature of α-amino acids and (IUPAC Commission on the Nomenclature of Organic Chemistry & IUPAC–IUB Commission on Biochemical Nomenclature) 1–16

Peptidoglycan, linkage between teichoic acid and, in the cell wall of a bacteriophage-resistant mutant of *Staphylococcus aureus* H (Heckels, J. E., Archibald, A. R. & Baddiley, J.) 637–647

Peroxidase, oxidation of ethyl formylphenylacetate by an enzyme system possessing activities of oxidase and, from *Datura innoxia* roots (Kalyanaraman, V. S., Mahadevan, S. & Kumar, S. A.) 565–576

Peroxidase, oxidation of reduced nicotinamide–adenine dinucleotide in the presence of ethyl formylphenylacetate by an enzyme system possessing activities of oxidase and, from *Datura innoxia* roots (Kalyanaraman, V. S., Kumar, S. A. & Mahadevan, S.) 577–584

Peroxidase, purification and properties of, from *Mycobacterium tuberculosis* H37Rv and its possible role in the mechanism of action of isonicotinic acid hydrazide on the organism (Gayathri Devi, B., Shaila, M. S., Ramakrishnan, T. & Gopinathan, K. P.) 187–197

pH, dependence on, of the hydrolysis of benzylpenicillin and ampicillin by *Bacillus cereus* 569/H β-lactamase I (Waley, S. G.) 547–551

Phenol sulphotransferase, properties of *Aspergillus oryzae* arylsulphatases, one possessing activity as (Burns, G. R. J. & Wynn, C. H.) 697–705

Phenolase, beet-leaf, kinetics of the hydroxylation of *p*-coumarate to caffeate by (McIntyre, R. J. & Vaughan, P. F. T.) 447–461

L-Phenylalanine ammonia-lyase, purification, subunit structure and kinetic properties of, from *Rhizoctonia solani* (Kalghatgi, K. K. & Subba Rao, P. V.) 65–72

L-Phenylalanine, kinetics of the formation of *trans*-cinnamate from, catalysed by *Rhizoctonia solani* L-phenylalanine ammonia-lyase (Kalghatgi, K. K. & Subba Rao, P. V.) 65–72

Phenylalanine residues, structural studies on three wheat-flour proteins that lack histidine residues and (Redman, D. G.) 725–732

Phenylalanyl-transfer ribonucleic acid, *N*-acetyl, interaction of thiol-containing analogues of puromycin with, on ribosomes from rat liver and *Escherichia coli* to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220

Phosphatase, alkaline, kidney, pig, mechanism of the hydrolysis of glycerol 2-phosphate by (Ahlers, J.) 535–546

Phosphatidylcholine, behaviour of mixed monolayers of fusogenic lipids and, and other phospholipids (Maggio, B. & Lucy, J. A.) 597–608

Phosphatidylinositol, comparison of the activities of rat brain phosphatidylinositiol phosphodiesterase towards, in dispersions and as monolayers at the air/water interface (Quinn, P. J. & Barenholz, Y. 199–208

Phosphatidylinositol phosphodiesterase, brain, rat, comparison of the activities of, towards phosphatidylinositol in dispersions and as monolayers at the air/water interface (Quinn, P. J. & Barenholz, Y.) 199–208

Phosphodiesterases, three, resolution, partial purification and properties of, from potato tuber (Ashton, A. R. & Polya, G. M.) 329–339

Phospholipase C, brain, rat, comparison of the activities of, towards phosphatidylinositol in dispersions and as monolayers at the air/water interace (Quinn, P. J. & Barenholz, Y.) 199–208

Phospholipids, behaviour of mixed monolayers of fusogenic lipids and (Maggio, B. & Lucy, J. A.) 597–608

Phosphorylase kinase, rabbit, phosphorylation of rabbit and ox heart-muscle troponin I by (Cole, H. A. & Perry, S. V.) 525–533

Pisum sativum, *see* Pea

Planorbis corneus, *see* Snail

Plasma membrane, *see* Membrane, plasma

Plasmacytoma cells, MPC-11, mouse, determination of polyadenylate-rich ribonucleic acid in the nucleus and cytoplasm of (Abraham, K. A. & Eikhom, T. S.) 669–674

Plastocyanin, *Anabaena variabilis*, amino acid sequence of, and its comparison with those of plastocyanins from *Chlorella fusca* and higher plants (Aitken, A.) 675–683

Polysaccharides, cell-wall, detection of acetyl groups in, in preparations from grasses and other higher plants (Bacon, J. S. D., Gordon, A. H., Morris, E. J. & Farmer, V. C.) 485–487

Polysaccharides, characterization of, obtained by mild acid hydrolysis of *Pseudomonas aeruginosa* N.C.T.C. 1999 lipopolysaccharide (Drewry, D. T., Symes, K. C., Gray, G. W. & Wilkinson, S. G.) 93–106

Potato (*Solanum tuberosum*) tuber, resolution, partial purification and properties of three cyclic nucleotide phosphodiesterases from (Ashton, A. R. & Polya, G. M.) 329–339

Potato (*Solanum tuberosum*) tuber, unidirectional inhibition and activation by *meso*-tartrate of 'malic' enzyme (EC 1.1.1.40) from (do Nascimento, K. H., Davies, D. D. & Patil, K. D.) 349–355

Protease, *see* Proteinase

Protein, fibrous (α-keratin), extraction and characterization of, from ox epidermis (Steinert, P. M.) 39–48

Protein, iron–sulphur, high-potential, *Chromatium* sp., X-ray photoelectron spectra of, and other iron–sulphur proteins (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474

Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, ox, phosphorylation of rabbit and ox heart-muscle troponin I by (Cole, H. A. & Perry, S. V.) 525–533

Protein kinase, adenosine 3′:5′-cyclic monophosphate-dependent, pitfall in the interpretation of data on the interaction of ligands with, and other proteins (Swillens, S. & Dumont, J. E.) 779–782

Protein, organization of chlorophyll in complexes with, from pea leaves (Scott, B. & Gregory, R. P. F.) 341–347

Protein–polysaccharide, *see* Proteoglycan

Protein, soluble, brain, rat, reciprocal exclusion by L-3,4-dihydroxyphenylalanine and L-tyrosine of their incorporation as single units into (Rodríguez, J. A., Barra, H. S., Arce, C. A., Hallak, M. E. & Caputto, R.) 115–121

Proteinase (acrosin), partial purification and properties of, from sheep spermatozoa (Brown, C. R., Andani, Z. & Hartree, E. F.) 133–146

Proteinase (acrosin), spermatozoal, sheep, determination of the operational molarity of, by fluorimetric titration with 4-methylumbelliferyl *p*-guanidinobenzoate (Brown, C. R., Andani, Z. & Hartree, E. F.) 147–154

Proteinase, *Armillaria mellea*, use of digestion with, having specificity for lysine residues in the establishment of the amino acid sequence of pig heart-muscle aspartate aminotransferase (Doonan, S., Doonan, H. J., Hanford, R., Vernon, C. A., Walker, J. M., Airoldi, L. P. da S., Bossa, F., Barra, D., Carloni, M., Fasella, P. & Riva, F.) 497–506

Proteins, analysis by two-dimensional gel electrophoresis of, of chick-embryo ribosome crystals (Dondi, P. G.) 475–476

Proteins, evidence against a proposed interaction of thionitrobenzoate with disulphide bonds of (Weitzman, P. D. J.) 281–283

Proteins, iron–sulphur, X-ray photoelectron spectra of (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474

Proteins, pitfall in the interpretation of data on the interaction of ligands with (Swillens, S. & Dumont, J. E.) 779–782

Proteins, reaction of benzylpenicillin with (Corran, P. H. & Waley, S. G.) 357–364

Proteins, reduction of disulphide bonds in, and protein mixed disulphides by a thioltransferase in a rat liver cytosol (Mannervik, B. & Axelsson, K.) 785–788

Proteins, wheat-flour, three, structural studies on, that lack phenylalanine and histidine residues (Redman, D. G.) 725–732

Proteoglycan, collagen-associated, isolation of, from ox nasal-septum cartilage and its preferential interaction with α_2 chains of calf skin type I collagen (Lee-Own, V. & Anderson, J. C.) 57–63

Proteoglycans, nature of the protein moieties of, of cartilage from pig larynx and ox nasal septum (Baxter, E. & Muir, H.) 657–668

Pseudomonas aeruginosa N.C.T.C. 1999, characterization of polysaccharide fractions obtained by mild acid hydrolysis of the lipopolysaccharide from (Drewry, D. T., Symes, K. C., Gray, G. W. & Wilkinson, S. G.) 93–106

Pseudomonas aeruginosa N.C.T.C. 8505, identification of 2-amino-2-deoxygalacturonic acid as a component of lipopolysaccharide from (Wilkinson, S. G. & Welbourn, A. P.) 783–784

Pseudomonas putida A.T.C.C. 17642, stereospecificity of nicotinamide–adenine dinucleotide-dependent dehydrogenases associated with the metabolism of *meso*-tartrate in (do Nascimento, K. H. & Davies, D. D.) 553–557

Puromycin, chemical synthesis of thiol-containing analogues of, and their interaction with *N*-acetylphenylalanyl-transfer ribonucleic acid on ribosomes from rat liver and *Escherichia coli* to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220

Pyridoxal 5′-phosphate, inactivation of horse liver alcohol dehydrogenase by reaction of, with the essential lysine residue (Chen, S.-S. & Engel, P. C.) 627–635

Pyridoxal 5′-phosphate, inactivation of pig muscle lactate dehydrogenase M_4 isoenzyme by reaction of, with an essential lysine residue (Chen, S.-S. & Engel, P. C.) 107–113

Pyridoxal 5′-phosphate, role of lysine residue 126 in ox liver glutamate dehydrogenase in the light of studies of the inhibition and inactivation of the enzyme by (Chen, S. -S. & Engel, P. C.) 619–626

Quinacillin, isolation of a specific covalent complex of *Staphylococcus aureus* PC1 penicillinase with (Virden, R., Bristow, A. F. & Pain, R. H.) 397–401

Red blood cells, *see* Erythrocytes

Rhizoctonia solani, purification, subunit structure and kinetic properties of L-phenylalanine ammonia-lyase from (Kalghatgi, K. K. & Subba Rao, P. V.) 65–72

Ribonucleic acid, messenger, activity of yeast mitochondrial ribonucleic acid as (Eggitt, M. J. & Scragg, A. H.) 507–512

Ribonucleic acid, mitochondrial, yeast, messenger activity of (Eggitt, M. J. & Scragg, A. H.) 507–512

Ribonucleic acid, polyadenylate-rich, determination of, in the nucleus and cytoplasm of mouse MPC-11 plasmacytoma cells (Abraham, K. A. & Eikhom, T. S.) 669–674

Ribonucleic acid polymerase, deoxyribonucleic acid-dependent, *Halobacterium cutirubrum* N.R.C. 34001, relationship between the deoxyribonucleic acid-bound and low-molecular-weight soluble forms of (Fitt, P. S., Peterkin, P. I. & Barua, N. N.) 719–724

Ribonucleic acid, ribosomal, 23S, *Agrobacterium tumefaciens*, presence of nicks in (Schuch, W. & Loening, U. E.) 17–22

Ribonucleic acid, transfer, *N*-acetylphenylalanyl-, interaction of thiol-containing analogues of puromycin with, on ribosomes from rat liver and *Escherichia coli* to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220

Ribonucleic acids, ribosomal, *Agrobacterium tumefaciens*, structure and biosynthesis of (Grienenberger, J. M. & Simon, D.) 23–30

Ribosomal ribonucleic acid, *see* Ribonucleic acid, ribosomal

Ribosome crystals, chick-embryo, analysis by two-dimensional gel electrophoresis of the proteins of (Dondi, P. G.) 475–476

Ribosomes, *Escherichia coli*, interaction of thiol-containing analogues of puromycin with *N*-acetylphenylalanyl-transfer ribonucleic acid on rat liver ribosomes and, to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220

Ribosomes, liver, rat, interaction of thiol-containing analogues of puromycin with *N*-acetylphenylalanyl-transfer ribonucleic acid on *Escherichia coli* ribosomes and, to form thioesters (Gooch, J. & Hawtrey, A. O.) 209–220

Rice (*Oriza sativa*) roots, effects of adenine nucleotides on the activity of adenosine triphosphate sulphurylase from (Onajobi, F. D.) 301–304

Roots, *Datura innoxia*, oxidation of ethyl formylphenylacetate by oxidase–peroxidase enzymes from (Kalyanaraman, V. S., Mahadevan, S. & Kumar, S. A.) 565–576

Roots, *Datura innoxia*, oxidation of reduced nicotinamide–adenine dinucleotide in the presence of ethyl formylphenylacetate by oxidase–peroxidase enzymes from (Kalyanaraman, V. S., Kumar, S. A. & Mahadevan, S.) 577–584

Roots, pea, glutamine and asparagine as nitrogen donors for the reductant-dependent biosynthesis of glutamate in (Miflin, B. J. & Lea, P. J) 403–409
Roots, rice, effects of adenine nucleotides on the activity of adenosine triphosphate sulphurylase from (Onajobi, F. D.) 301–304

Saccharide, stopped-flow fluorescence studies on the binding of, to hen's-egg-white lysozyme (Halford, S. E.) 411–422
Saccharomyces cerevisiae, messenger activity of mitochondrial ribonucleic acid from (Eggitt, M. J. & Scragg, A. H.) 507–512
Saliva, parotid-gland, human, identification in, of a peptide particularly active in enhancing the glycolytic activity of salivary micro-organisms (Holbrook, I. B. & Molan, P. C.) 489–492
Salivary gland, parotid, human, identification in saliva from, of a peptide particularly active in enhancing the glycolytic activity of salivary micro-organisms (Holbrook, I. B. & Molan, P. C.) 489–492
Satellite deoxyribonucleic acid, *see* Deoxyribonucleic acid, satellite
Sea-cucumber (*Holothuria forskali*), effects of anions on the activity of arginine kinases from lobster (*Homarus vulgaris*) and (Anosike, E. O. & Watts, D. C.) 387–395
Serum apolipoproteins, *see* Apolipoproteins, serum
Skeletal muscle, *see* Muscle, skeletal
Skin, calf, isolation of a collagen-associated proteoglycan from ox nasal-septum cartilage and its preferential interaction with α_2 chains of type I collagen from (Lee-Own, V. & Anderson, J. C.) 57–63
Skin epithelial cells, human, photochemical binding of anthracene to deoxyribonucleic acid of monkey kidney epithelial cells and, in tissue culture (Blackburn, G. M. & Taussig, P. E.) 289–291
Skin, ox, extraction and characterization of epidermal α-keratin from (Steinert, P. M.) 39–48
Skin, ox, mechanism of the stabilization of the reducible intermediate cross-links of collagens from, and other tissues (Robins, S. P. & Bailey, A. J.) 381–385
Snail (*Planorbis corneus*), physicochemical properties of erythrocruorin from (Wood, E. J. & Mosby, L. J.) 437–445
Solanum tuberosum, *see* Potato
Sorbitol 6-phosphate dehydrogenase, nicotinamide–adenine dinucleotide-dependent, *Aerobacter aerogenes* N.C.I.B. 8017, stereospecificity of, and mannitol 1-phosphate dehydrogenase (do Nascimento, K. H. & Davies, D. D.) 553–557
Spectrophotometer, stopped-flow, rapid-scanning, double-beam construction of, and its use for the measurement of rapid spectral changes (Hollaway, M. R. & White, H. A.) 221–231
Spermatozoa, sheep, determination of the operational molarity of acrosin from, by fluorimetric titration with 4-methylumbelliferyl *p*-guanidinobenzoate (Brown, C. R., Andani, Z. & Hartree, E. F.) 147–154
Spermatozoa, sheep, partial purification and properties of acrosin from (Brown, C. R., Andani, Z. & Hartree, E. F.) 133–146
Spinacea oleracea, *see* Spinach
Spinach-beet, *see* Beet
Spinach (*Spinacea oleracea*), X-ray photoelectron spectra of ferredoxin from and of other iron–sulphur proteins (Andrews, P. T., Johnson, C. E., Wallbank, B., Cammack, R., Hall, D. O. & Rao, K. K.) 471–474
Staphylococcus aureus H, linkage between teichoic acid and peptidoglycan in the cell wall of a bacteriophage-resistant mutant of (Heckels, J. E., Archibald, A. R. & Baddiley, J.) 637–647
Staphylococcus aureus PC1, isolation of a specific covalent complex of penicillinase from, with quinacillin (Virden, R., Bristow, A. F. & Pain, R. H.) 397–401
Steoroyl-coenzyme A, regulatory effects of, and other fatty acyl-coenzyme A species on the activity of pig brain and kidney phosphate-activated glutaminase *in vitro* (Kvamme, E. & Torgner, I. Aa) 83–91
Sulphate esters, secondary alkyl, activity towards, of secondary alkylsulphatases in a strain of *Comamonas terrigena* (Fitzgerald, J. W.) 477–480
N- and *O*-Sulphotransferases, solubilization and partial characterization of, involved in the biosynthesis of heparin from mouse mastocytoma microsomal fraction (Jansson, L., Höök, M., Wasteson, Å. & Lindahl, U.) 49–55
Supernatant fraction, *see* Cytosol

meso-Tartrate, stereospecificity of nicotinamide–adenine dinucleotide-dependent dehydrogenases associated with the metabolism of, in *Pseudomonas putida* A.T.C.C. 17642 (do Nascimento, K. H. & Davies, D. D.) 553–557
meso-Tartrate, unidirectional inhibition and activation by, of potato-tuber 'malic' enzyme (EC 1.1.1.40) (do Nascimento, K. H., Davies, D. D. & Patil, K. D.) 349–355
Teichoic acid, *Lactobacillus buchneri* N.C.I.B. 8007, interaction of magnesium ions with (Lambert, P. A., Hancock, I. C. & Baddiley, J.) 519–524
Teichoic acid, linkage between peptidoglycan and, in the cell wall of a bacteriophage-resistant mutant of *Staphylococcus aureus* H (Heckels, J. E., Archibald, A. R. & Baddiley, J.) 637–647
Temperature, method for the investigation of the effect of, on the 695 nm absorption band of immobilized horse heart cytochrome *c* (Moore, T. A. & Greenwood, C.) 169–177
Tendon, ox, mechanism of the stabilization of the reducible intermediate cross-links of collagens from, and other tissues (Robins, S. P. & Bailey, A. J.) 381–385
Testis, ox, properties of follicle-stimulating-hormone receptor of plasma membrane from (Cheng, K.-W.) 123–132
Thioltransferase, cytosolic, liver, rat, reduction of disulphide bonds in proteins and protein mixed disulphides by (Mannervik, B. & Axelsson, K.) 785–788
Thionitrobenzoate, evidence against a proposed interaction of, with disulphide bonds of proteins (Weitzman, P. D. J.) 281–283
Transfer ribonucleic acid, *see* Ribonucleic acid, transfer
Transferrin, ovo-, chicken, electron-paramagnetic-resonance spectroscopy of iron-binding fragments of (Butterworth, R. M., Gibson, J. F. & Williams, J.) 559–563

Transferrin, ovo-, chicken, preparation and properties of iron-binding fragments from the *C*-terminal region of (Williams, J.) 237–244
Triticum aestivum, *see* Wheat
Troponin I, heart-muscle, rabbit and ox, phosphorylation of (Cole, H. A. & Perry, S. V.) 525–533
Troponin I, skeletal-muscle, rabbit, amino acid sequence of (Wilkinson, J. M. & Grand, R. J. A.) 493–496
Trypsin, ox, determination of the operational molarity of sheep spermatozoal acrosin and, by fluorimetric titration with 4-methylumbelliferyl *p*-guanidinobenzoate (Brown, C. R., Andani, Z. & Hartree, E. F.) 147–154
Tryptophan residue, kinetic studies on horse heart cytochrome *c* *N*-formylated at (Brittain, T. & Greenwood, C.) 713–717
Tryptophan residue, kinetic studies on horse heart cytochrome *c* modified by reaction at, with 2-hydroxy-5-nitrobenzyl bromide (Brittain, T. & Greenwood, C.) 179–185
Tuber, potato, resolution, partial purification and properties of three cyclic nucleotide phosphodiesterases from (Ashton, A. R. & Polya, G. M.) 329–339
Tuber, potato, unidirectional inhibition and activation by *meso*-tartrate of 'malic' enzyme (EC 1.1.1.40) from (do Nascimento, K. H., Davies, D. D. & Patil, K. D.) 349–355
Tubulin, reciprocal exclusion by 3,4-dihydroxyphenylalanine and L-tyrosine of their incorporation as single units into a rat brain soluble protein tentatively identified as (Rodríguez, J. A., Barra, H. S., Arce, C. A., Hallak, M. E. & Caputto, R.) 115–121
Tyramine, effect of, on the activity of *Aspergillus oryzae* arysulphatases (Burns, G. R. J. & Wynn, C. H.) 697–705
L-Tyrosine, kinetics of the formation of *p*-coumarate from, catalysed by *Rhizoctonia solani* L-phenylalanine ammonia-lyase (Kalghatgi, K. K. & Subba Rao, P. V.) 65–72
L-Tyrosine, reciprocal exclusion by L-3,4-dihydroxyphenylalanine and, of their incorporation as single units into a rat brain soluble protein (Rodríguez, J. A., Barra, H. S., Arce, C. A., Hallak, M. E. & Caputto, R.) 115–121

Wall, cell, *see* Cell wall
Wheat (*Triticum aestivum*) flour, structural studies on three proteins from, that lack phenylalanine and histidine residues (Redman, D. G.) 725–732

Xenopus laevis, messenger activity of yeast mitochondrial ribonucleic acid injected into oocytes of (Eggitt, M. J. & Scragg, A. H.) 507–512
Xylitol dehydrogenase, liver, guinea-pig, nicotinamide-adenine dinucleotide-dependent, stereospecificity of, and of other dehydrogenases (do Nascimento, K. H. & Davies, D. D.) 553–557

Yeast (*Saccharomyces cerevisae*), messenger activity of mitochondrial ribonucleic acid from (Eggitt, M. J. & Scragg, A. H.) 507–512

Zinc ions, induction of metallothionein in rat liver by the injection of, and by the restriction of food intake (Bremner, I. & Davies, N. T.) 733–738